

REFRACTION OF THE EYE

MORTON.

FOURTH EDITION



4/6

X Feb. 12

R37799

REFRACTION OF THE EYE



REFRACTION OF THE EYE

ITS

DIAGNOSIS

AND THE

CORRECTION OF ITS ERRORS

BY

A. STANFORD MORTON, M.B., F.R.C.S. ENG.

ASSISTANT SURGEON TO THE MOORFIELDS OPHTHALMIC HOSPITAL; OPHTHALMIC
SURGEON TO THE GREAT NORTHERN CENTRAL HOSPITAL; LATE SURGEON
TO THE ROYAL SOUTH LONDON OPHTHALMIC HOSPITAL



FOURTH EDITION

LONDON

H. K. LEWIS, 136 GOWER STREET, W.C.

1891

PRINTED BY
H. K. LEWIS, 136 GOWER STREET
LONDON, W.C.

PREFACE TO THE FOURTH EDITION.

THE subject matter of this edition has been revised, and partly rewritten in such a way as to make it more practically useful to the student and practitioner.

Specimens of test-types are appended as in the previous edition.

WEYMOUTH STREET, W.

October, 1891.

PREFACE TO THE FIRST EDITION.

THESE notes are published with a view of enabling Practitioners to diagnose, and correctly estimate the value of, the phenomena indicating the state of a patient's refraction. They are intended to furnish a basis for observation; and it is hoped that they will make evident the necessity which exists for *personally working out* a large number of refraction cases in order to acquire anything like proficiency in prescribing correct glasses.

To those friends who have aided me by their suggestions I would take this opportunity of expressing my best thanks.

March, 1881.

TABLE OF CONTENTS.

	PAGE
INTRODUCTION	I
CHAPTER I.	
Action of Rays and Lenses	3
CHAPTER II.	
Definitions	6
CHAPTER III.	
Old and New Systems of Lenses—Table of Comparison . . .	10
CHAPTER IV.	
Accommodation	13
CHAPTER V.	
Conditions necessary for clear perception of a line . . .	16
CHAPTER VI.	
Description of Test-types—Mode of use	18
CHAPTER VII.	
Indications afforded by use of Test-types	21

CHAPTER VIII.

	PAGE
Indications afforded by Mirror alone—Movements of Image of disc in Emmetropia, Myopia, Hypermetropia and Astigmatism— Explanation of same	23

CHAPTER IX.

Retinoscopy	30
-----------------------	----

CHAPTER X.

Indications afforded by mirror and lens—Change in size of Image of disc in Emmetropia, Myopia, Hypermetropia and Astig- matism—Explanation	46
--	----

CHAPTER XI.

Estimation of Refraction by Ophthalmoscope—Condition of Ob- server's own Refraction—Tabular view of different States of Refraction	51
--	----

CHAPTER XII.

Test-types and glasses—Prescribing Spectacles in Hypermetropia— Myopia — Astigmatism — Presbyopia — Table of glasses for Presbyopia—Aphakia	56
---	----

CHAPTER XIII.

Action of Atropine—Homatropine—Homatropine and Cocaine— Eserine—Calculation for same	66
---	----

CHAPTER XIV.

Amblyopia—Malingering and its detection	70
---	----

REFRACTION OF THE EYE.

INTRODUCTION.

THESE pages have been written in the hope that they may be of service to beginners, and to Physicians and general Practitioners, who, systematically using the Ophthalmoscope in their investigation of disease, must avail themselves of the information thereby afforded regarding the patient's refraction, in order that due allowance may be made for any error which exists. If patients can assist by their answers, much valuable information is afforded by the use of test-types and glasses, a description of which, with the information to be derived therefrom, has been therefore introduced.

Information is given on the subject of the glasses required in the more *ordinary* cases, but for a knowledge of the various diseases which accompany and complicate many of the errors of refraction, the reader is referred to the works already published on that subject, to the study of which these notes are intended to prepare the way.

So essential is a constant remembrance of some of the facts hereafter mentioned, that, at the risk of tautology, they have been kept constantly before the reader. The plan which has been found so useful in other branches of medical study has been adopted here, viz., that of working out from the symptoms the nature of the disease in prefer-

ence to the method of naming the disease and then describing the symptoms accompanying it.

In making an estimation of the refractive character of an eye, it is especially necessary to proceed *systematically*. On a patient complaining of "bad sight" the vision of each eye must always be tested separately, and we should examine him somewhat in the following manner:—

1. Listen carefully to the nature of his complaint.
2. Test the near and distant vision *without* glasses.
3. Estimate the refractive character of the eye by either
 - (a) Ophthalmoscopy,
 - (b) Retinoscopy ; .

both of which are *objective* methods.

4. Examine the eye with an ophthalmoscope in order to ascertain whether any lesion other than that of Refraction exists.

5. Test the distant and near vision with the glasses for whose use the previous examination has indicated the necessity.

Two methods of notation are at present in vogue (*vide* Chap. III.). The new system of numeration is the one which is now almost universally employed, and it has been adopted in the text. The *approximately* corresponding measurements in English inches have, however, been given in brackets for the convenience of those who prefer the old method.

Only so much ophthalmoscopy has been introduced as was necessary to explain the phenomena of refraction.


CHAPTER I.

ACTION OF RAYS AND LENSES.

a. RAYS of light issuing from every point of any object diverge in *all* directions. The nearer an eye is to such an object the more of its *diverging* rays does it intercept, and *vice versa*, the further the eye is removed, the fewer are the divergent rays which reach it: *i.e.*, the more parallel are the rays which enter it. Only from objects at an infinite distance do we receive absolutely parallel rays.

For practical purposes, however, we may *consider* those rays to be parallel which proceed from an object situated at a distance of 6 metres (20 ft.) or more.

FIG. 1.




b. Parallel rays (Fig. 1, Aa, Bb) passing through a convex spherical lens come to a focus at a point, F, on the other side of the lens. This point is called the "principal focus" of the lens, and the distance from this point to the centre of the lens is named "the focal distance" or "focal length" of the lens. Conversely, rays proceeding from a point situated at the principal focus of the lens emerge parallel to each other.

c. *Diverging rays* (Fig. 1, *Ma*, *Mb*) after traversing such a lens unite at a point, *K*, which is situated further from the lens than the principal focus. Conversely, rays proceeding from a point further from the lens than its principal focus, after passing through it, emerge as convergent rays.

d. *Converging rays* (Fig. 1, *Ca*, *Cb*) after traversing a convex lens meet at a point, *H*, which is nearer to the lens than its principal focus. Conversely, rays from a point situated nearer to the lens than the principal focus, are, on emerging from the lens, still divergent, though less so than before passing through it.

FIG. 2.



e. Parallel rays (Fig. 2, *Aa*, *Bb*) after traversing a concave spherical lens diverge (*aC*, *bC*) as if proceeding from a point, *F*, "the virtual focus," situated on the *same* side of the lens as that from which the parallel rays proceed. The distance of this point from the lens is the "negative focal length" of the lens. Conversely, rays which converge as if proceeding to this point are rendered parallel by passing through it.

f. A cylindrical lens influences the course taken by rays of light in the same manner as a spherical, but acts most on those proceeding in a direction at *right angles* to its axis.

g. By the *Refraction* of a lens is meant the property

which it possesses of altering the direction of rays traversing it.

h. The refracting *power* of a lens is indicated by the distance from it of the point at which parallel rays traversing it meet, *i.e.* by its focal distance. The greater the focal distance the less is its refracting power and *vice versa*.


CHAPTER II.

DEFINITIONS.

DEF. 1. The refracting surfaces of the eye are, according to Donders—*a*, the anterior surface of the cornea; *b*, the anterior surface of the lens; *c*, the anterior surface of the vitreous. The transparent media together form the “*Dioptric system*” of the eye, which influences the course taken by rays of light traversing it in a manner very similar to a biconvex lens.

DEF. 2. “*Refraction of the eye*” means the effect which, by reason of its form and structure, this organ produces upon rays of light entering it when the action of the accommodation apparatus is completely suspended.

FIG. 3.




DEF. 3. In *Emmetropia* (Fig. 3) parallel rays, *Aa*, *Bb*, come to a focus upon the retina, *F*, when there is no accommodation. Conversely, rays proceeding from points in the retina emerge from the eye, parallel to one another.

DEF. 4. An *Ametropic* eye is one in which, when there is no accommodation, parallel rays do *not* come to a focus upon the retina. Such an eye may be either Myopic, Hypermetropic or Astigmatic.


DEF. 5. A *Myopic* eye (Fig. 4) is that in which, when there is no accommodation, parallel rays come to a focus in *front* of the retina, as at F. Conversely, rays issuing from the retina, R, which is behind the principal focus of the

FIG. 4.



dioptric system, emerge as convergent rays (Chap. I., *c*) which meet at a point M, in front of the eye. Rays proceeding from this point (called the "far point") would therefore be focussed on the retina.

FIG. 5.



DEF. 6. A *Hypermetropic* eye (Fig. 5) is that in which, when there is no accommodation, parallel rays, (*Aa*, *Bb*) are so changed in direction that they tend to come to a focus at F, *behind* the retina. Conversely, rays proceeding as if from F, would emerge from the eye parallel to one another. Rays, therefore, from R, which is situated *within* the focus of the dioptric system (*vide* Def. 1) emerge from the eye diverging (Chap. I., *d*) as if they were proceeding from a point G.

The distance of this point behind the front part (nodal point) of the eye will, in any case, be equal to the focal length of the convex lens which, when held close to

the eye with its accommodation suspended, would bring parallel rays to a focus on the retina. For since the rays from the retina, on emerging from the eye, diverge as if they issued from G, it follows that rays converging, as if proceeding to G, would meet on the retina at R. Consequently, the convex lens which, with the aid of the dioptric system of the eye, brings parallel rays to a focus at R, would without the aid of that system bring them to a focus at G. This latter point is therefore at the principal focus of the lens (*vide* Chap. I., *b*).

DEF. 7. A *regularly* astigmatic eye is that in which there is a difference in the degree of refraction in different meridians; the two *principal* meridians (that is the meridians of greatest and least error) being always at *right angles* to each other.

“Regular Astigmatism” may exist in five different forms:—

- a.* Simple myopic = one principal meridian emmetropic, the other myopic.
- b.* Simple hypermetropic = one principal meridian emmetropic, the other hypermetropic.
- c.* Compound myopic = both principal meridians myopic, one more than the other.
- d.* Compound hypermetropic = both principal meridians hypermetropic, one more than the other.
- e.* Mixed = one principal meridian myopic, the other hypermetropic.

DEF. 8. An *Irregularly Astigmatic* eye is one in which there are different degrees of refraction in different parts of the various meridians.

DEF. 9. A *Presbyopic* eye is that in which the power of accommodating for near objects is lost, owing to physiological changes produced by advancing age, and there is, consequently, an inability to define small objects situated near to the eye (*vide* Presbyopia, Chap. XII.). It may co-exist with any of the above named conditions.

DEF. 10. *Anisometropia* is a term used to denote that the refractive condition of one eye differs from that of the other.

From the statements in this and the preceding chapter the following deductions may be made.

Deduction 1. In order that *parallel* rays may be brought to a focus on the retina of a myopic eye they must be made to diverge as if they proceeded from the "far-point" (Def. 5, p. 7). This is effected by placing before the eye and close to it the concave lens whose negative focal length equals the distance of this point from the eye. (Chap. I., *e*). The amount of the Myopia is said to be equal to the refracting power (Chap. I., *h*) of this lens.

Deduction 2. In order that *parallel* rays may be brought to a focus on the retina of a hypermetropic eye at rest, they must be made to converge as if they proceeded to the point behind the eye from which retinal rays appeared to issue (p. 7, fig. 5, G). This is effected by placing before and close to the eye a biconvex lens whose focal length is equal to the distance of that point from the front of the eye. The *amount* of the Hypermetropia is said to be equal to the refracting power of this lens.

Deduction 3. In regular astigmatism one principal meridian is corrected by a cylindrical lens. Correction of both requires a bi- or spherico-cylindrical lens.

CHAPTER III.

OLD AND NEW SYSTEMS OF MEASUREMENT.

IN the old system, the unit of measurement was a lens whose focal length or refracting power was 1 inch.

Such a strong lens did not, however, exist in the trial cases, and consequently the numbers of the lenses were expressed by some *fraction* ($\frac{1}{2}$, $\frac{1}{40}$, etc.) indicating how much their refracting power was less than the lens whose focal distance was 1 inch.

The denominator of the fraction indicated, in inches, the focal distance of any given lens.

It will thus be seen that the lenses whose focal distances were 6 inches and 12 inches, had respectively $\frac{1}{6}$ and $\frac{1}{12}$ the refracting power of the lens whose focal distance was 1 inch. The intervals between the lenses were irregular and the difference between the refracting powers, of any two lenses, had to be calculated by means of *fractions*.

In the new or "Dioptric" system, the unit of measurement is a lens, whose focal distance or refracting power is 1 Metre or "Dioptre," = 1 D. If two such lenses were placed together they would have twice the refracting power of this unit with half its focal distance.

A lens, therefore, having a refracting power equal to two such units, is expressed as 2 D. and has a focal distance of 50 cm. In the same way a lens equal in refracting power

to ten such units = 10 D., and its focal distance is one tenth of that of the unit (= 10 cm.). Further, a lens whose refracting power is only half that of the unit = .5 D. and has twice the focal distance of the unit, viz., 200 cm.

The Dioptric system of notation, has now almost universally superseded the other system, because the intervals between the lenses are regular and the calculations are made in *whole numbers* instead of fractions.

The focal length of any lens can be instantly estimated. Thus, a division of the number of the lens in dioptries (D.) into 100 gives the focal length of the lens in centimetres. Conversely the division of the focal length (in centimetres) of any lens into 100, gives the strength of the lens in dioptries.

In order to ascertain the corresponding numbers in the two systems for any lens, we must remember that 1 dioptre (or metre) = 37 Paris inches and (39.4) nearly 40 English inches, and that, therefore, the lens with 1 D. focal length corresponds to the lens in the old series whose focal length is 40 ins. ($\frac{1}{40}$); 2 D. to a lens with double this refracting power or $\frac{2}{40}$ ($\frac{1}{20}$) of the old, and so on.

Where d , then, represents the number of dioptries, and a the number of inches, we can ascertain (i) the lens in the old series which corresponds to one of the new, by the formula $\frac{d}{40} = \frac{1}{a}$; and (ii) the number of a lens in the new system corresponding to one in the old by the formula $d = \frac{40}{a}$.

The following table gives the lens which, in an ordinary

French or English case, will correspond *most* nearly with any dioptré.

For greater accuracy in any special case, the reader must calculate according to the formulæ given above.

TABLE I.

D	Focal length in English inches.	Focal length in Paris inches.	Focal length in centimetres.	D	Focal length in English inches.	Focal length in Paris inches.	Focal length in centimetres.
0.25	160	144	400	5.0	8	7	20
0.50	80	72	200	5.50	$7\frac{1}{2}$	$6\frac{1}{2}$	18
0.75	50	50	133	6.0	7	6	16
1.0	40	36	100	7.0	6	5	14
1.25	30	30	80	8.0	5	$4\frac{1}{2}$	12.5
1.50	24	24	66	9.0	$4\frac{1}{2}$	4	11
1.75	22	20	57	10.0	4	$3\frac{3}{4}$	10
2.0	20	18	50	11.0	$3\frac{1}{2}$	$3\frac{1}{2}$	9
2.25	18	16	44	12.0	$3\frac{1}{4}$	3	8.3
2.50	16	14	40	13.0	3	$2\frac{3}{4}$	7.6
2.75	14	13	36	14.0	$2\frac{3}{4}$	$2\frac{1}{2}$	7.1
3.0	12	12	33	15.0	$2\frac{3}{4}$	$2\frac{1}{2}$	6.6
3.50	11	10	28	16.0	$2\frac{1}{2}$	$2\frac{1}{4}$	6.2
4.0	10	9	25	18.0	$2\frac{1}{4}$	2	5.5
4.50	9	8	22	20.0	2	$1\frac{3}{4}$	5

CHAPTER IV:

ACCOMMODATION.

DEF. II. By "the accommodation of the eye" is meant the power which we possess of bringing to a focus on the retina, rays proceeding from objects situated at various distances. These distances range between the "far point" and the "near point." The "far point" (*punctum remotum*, p.r.) is the point for which the eye is adjusted when the accommodation is totally suspended, and the "near point" (*punctum proximum*, p.p.) is the point for which the eye is adjusted when the whole of the accommodative power is called into activity. The amount of accommodation thus exercised by an eye in passing from a condition of extreme relaxation to a condition of maximum tension is called the *Range* or *Amplitude* (Donders) of accommodation. This amplitude may be represented by the convex lens which, with completely suspended accommodation, would bring rays proceeding from the "near point" to a focus on the retina.

Now a biconvex lens whose focal length equals the distance of the near point from the eye will, when placed close to an eye, render *parallel* the rays entering that organ from its near point. (Vide Chap. I., b).

Since in *Emmetropia* (E.), when the eye is at rest, *parallel* rays come to a focus on the retina (Def. 3, p. 6), such a lens represents in E. the amplitude of accommodation.

But in *Hypermetropia* (H.), when the eye is at rest, *parallel* rays do *not* come to a focus on the retina without the aid of a convex lens (Chap. II., Ded. 2). In order to estimate the amplitude of accommodation in this case the strength of the lens necessary to render the parallel rays sufficiently convergent, must therefore be *added* to that of the lens whose focal length equals the distance of the near point from the eye. Thus, if the H. = 3 D. ($\frac{1}{12}$), and the near point = 14 cm. (6 ins.), then, to the lens whose focal length = 14 cm., viz. 7 D. ($\frac{1}{6}$), we add that which brings parallel rays to a focus on the retina, viz. 3 D. ($\frac{1}{12}$). The amplitude therefore will be 7 D. + 3 D. = 10 D. ($\frac{1}{6} + \frac{1}{12} = \frac{1}{4}$).

From this it will be seen that, with the *same* amplitude of accommodation, the near point is *further* from the eye in H. than in E.

In *Myopia* (M.), since a concave lens is necessary to effect the focussing of parallel rays on the retina (Chap. II., Ded. 1), it is evident that in order to estimate the amplitude of accommodation, the strength of the lens requisite to render parallel rays sufficiently divergent to come to a focus on the retina, must be *deducted* from that of the lens whose focal length equals the distance of the near point from the eye. Thus, if the M. = 3 D. ($\frac{1}{12}$), (i.e. p.r. = 33.3 cm.), and the near point = 7.6 cm. (3 ins.), then, from the lens whose focal length = 7.6 cm., viz. 13 D., we must *deduct* that which brings parallel rays to a focus on the retina, viz. 3 D. ($\frac{1}{12}$). The amplitude, therefore will be 13 D. — 3 D. = 10 D. ($\frac{1}{3} - \frac{1}{12} = \frac{1}{4}$). From this we see that, with the *same* amplitude of accommodation, the near point is *closer* to the eye in M. than in E.

It is further evident that, if the "near point" had been the *same* for each of the foregoing examples, the amplitude would have been greater in H., and less in M., than in E.

As age advances the near point recedes further from the eye, so that in *Presbyopia* there is an *absolute* diminution of the amplitude. The amount of the diminution varies with the age of the individual. If the near point recedes until it reaches the far point, the accommodation of course becomes *nil*.

CHAPTER V.

PERCEPTION OF A LINE.

THE distinctness with which a line is visible depends upon the sharp and well-defined perception of its *margins*; if these are indistinct the line appears hazy. A line may be taken as made up of an infinite number of elements or points, from each of which rays issue in all directions. To gain a distinct image of any line, it is necessary that the rays from these points, which emerge in planes at *right angles* to its long axis, should be brought to a focus at points on corresponding planes on the retina, otherwise circles of diffusion are formed in this *transverse* direction of the line, which overlap each other, and finally, by projecting beyond its margins, give to it an ill-defined and blurred outline, so that no distinct perception of it is obtained. If from these same points, the rays emerging in planes *parallel* to the long axis overlap each other, it is only at the two extremities of the line that, by projecting, they cause any blurring. The margins of the line, thus not being in any way affected, there is no interference with the outline, and a clear image is formed on the retina.

If then a patient with his accommodation suspended, and who is emmetropic in one meridian, and myopic or hypermetropic in the other, be placed at 6 m. (20 ft.) from radiating lines of equal definition, he will see most *distinctly* that line which runs at *right angles* to his *Emmetropic* meri-

dian. Rays from points in the transverse planes of this line will, by passing through the emmetropic meridian, come to a focus on his retina, giving a distinct, well-defined image of the margins, and hence a clear perception of the line. The line *parallel* to the emmetropic meridian will, at the same time, be the most *indistinct*. Rays from *its* transverse planes pass through the myopic or hypermetropic meridian. They thus come to a focus, in the former case, in front of, and in the latter behind, the retina, producing circles of diffusion in these planes, and a consequent blurring of the margins of the line, so that no distinct image is obtained.

RULE I.—We have then the rule that in *simple* astigmatism, the patient, at 6 m., sees *most distinctly* the line *parallel* to the plane of his *error of refraction*.

It follows also, that a patient with either compound or mixed astigmatism, will not see *any* line distinctly at 6 m. when his accommodation is suspended.

CHAPTER VI.

DESCRIPTION OF TEST-TYPES: METHOD OF EMPLOYING THEM.

TEST-TYPES are divided into—*a*, those adapted for ascertaining *near* vision; and—*b*, those adapted for ascertaining *distant* vision. In using them we are obviously dependent on the answers given by the patient. For children, illiterate adults, and impostors, they are therefore an inferior means of diagnosing and estimating the refractive condition of an eye.

a. TEST-TYPES FOR NEAR VISION.

Those generally in use for this purpose are Jaeger's or Snellen's. The latter are so graduated that each should be read *as far off* as the distance for which it is marked. The smallest should be seen at a distance of 50 cm. (18 ins.) and the largest at 4 m. (12 ft.). These types are given into the patient's hand, and we then note the smallest he can read, and also the nearest and the farthest points at which it is distinctly visible. The small types are chiefly useful in testing the accommodation, but they also afford an indication of the presence and amount of myopia (*vide* Chap. XII., Myopia).

The wire optometer is perhaps the best means of ascertaining the position of the near point.

δ. TEST-TYPES FOR DISTANT VISION.

Snellen's types for this purpose are so graduated that each should be distinctly legible *as far off* as the distance for which it is marked. The largest should be seen at 60 metres, and each succeeding line at 36, 24, 18, 12, 9, and 6 metres respectively. The corresponding lines in the old system were of such a size as to be legible at 200, 100, 70, 50, 40, 30, and 20 feet respectively. Those who were accustomed to the old system, but who now use the new metrical type, still speak of the 2nd, 3rd, and 4th lines as if they were to be read at 100, 70, and 50 feet respectively, and they are therefore thus referred to in these pages, though these distances do not absolutely correspond to the distances at which these lines are meant to be read. The distances at which the rest of the lines are to be read correspond exactly in the two systems. The size of these types is such that any one with fairly good vision can read them, and consequently many young Emmetropes can see them clearly at a distance greater than that for which they are marked.

In testing with these types, we place the patient at such a distance that rays issuing from them reach his eye as parallel rays. For *practical* purposes this is obtained at a distance of 6 m. (20 ft.) (*vide* Chap. I., a) from which point the lowest line should be read by the normal eye without accommodation: and in these pages this is the distance at which the patient is always assumed to stand. By many,

however, 5 m. (16 ft.) is considered to be a sufficient distance. For testing astigmatism some use a fan of radiating lines all of the same magnitude and definition. These should all appear equally distinct to the normal eye at 6 m. In noting the vision (V.) we employ a fraction whose numerator denotes the distance at which the patient stands, and whose denominator indicates the lowest line which he can read. The number of the line is designated by the distance in metres (or feet), at which it *should* be legible. Thus normal $V. = \frac{6}{6} (\frac{20}{20})$ or 1; but if at 6 m. (20 ft.) the patient read only the line which *should* be read as far off as 12 m. (40 ft.) we say $V. = \frac{6}{12} (\frac{20}{40})$ or $\frac{1}{2}$. Seeing that our object in testing refraction is always to have the accommodation suspended, we should not place the patient nearer the types than 6 m. If it be necessary, however, in defective V. *from other causes*, we may allow him to approach the types till he can read the largest; if this be at a distance of, say 2 m. (6 ft.), then $V. = \frac{2}{60} (\frac{6}{200})$.

Dr. Snellen's test types may be obtained from Messrs. Williams and Norgate, Henrietta Street, Covent Garden, but some types corresponding to those arranged by him have by his kind permission been introduced at the end of this book for purposes of illustration. The number over each type indicates in metres and feet the distance at which it should be legible, and, in the types for testing near vision, the figure in the right hand corner denotes the approximately corresponding number of Jaeger's type.

CHAPTER VII.

INDICATIONS AFFORDED BY USE OF TEST-TYPES.

NOTE.—In the following section the patient is tested *without* glasses, then :—

Indication 1. If a patient read Jaeger 1 (J. 1) or Snellen 1 (Sn. 1) with a good range, and read also $\frac{6}{8}$ ($\frac{20}{20}$) perfectly, he is probably *emmetropic*. He cannot be myopic though he may be hypermetropic: for a hypermetropic patient with active accommodation could accomplish this.

Indication 2. If a patient *over 40 years of age* read only the larger J. or Sn. types (or perhaps even the smaller) but only on condition that he holds them at a considerable distance; while at the same time he can read $\frac{6}{8}$ ($\frac{20}{20}$) perfectly, he is simply presbyopic.

Indication 3. If a patient must hold the types *close* to his eye but can then read even the *finest* though he cannot see $\frac{6}{80}$ ($\frac{20}{600}$) he is *myopic*.

NOTE.—A patient may read Sn. 1 as far as 50 cm. (18 ins.) or even Sn. 2 as far as 60 cm. (2 ft.) together with some of the larger distance types such as $\frac{6}{36}$ ($\frac{20}{100}$). In this case he is *very slightly myopic*. N.B. Since such a patient can read Sn. 1 up to within a short distance from his eye he is thus easily distinguished from the following.

Indication 4. If a patient read only the larger series of the small types, and the smaller series not at all, or only

very imperfectly : while at the same time the distant vision is very defective, we suspect either some form of *astigmatism* or *hypermetropia without accommodation*.

In a patient *under* 40 years of age it is probably the former, but in one *over* that age it may be *either*, though it is frequently only the latter condition which exists.

Indication 5. If a patient *under* 40 or 45 can read only such large types as Sn. for 4 m. (J. 16) while he can read $\frac{6}{6}$ ($\frac{20}{20}$) he has *paralysis of accommodation*. This may be proved by giving him a strong convex lens such as +3 D. (12 ins.) when he will read Sn. 1, or J. 1, at the focal distance of the lens. (Chap. IV., p. 13).

Indication 6. If a patient, whose accommodation is suspended, see, quite distinctly, *one only* of the radiating lines at 6 m., he is *emmetropic* only in the meridian at *right angles* to this line, and either hypermetropic or myopic in the other. (Chap. V., p. 16).

CHAPTER VIII.

INDICATIONS AFFORDED BY MEANS OF MIRROR ALONE.

"DIRECT METHOD."

IN order that the indications of the refraction, afforded by this method, may be realised, the observer, if he be not emmetropic, must correct any error of his own refraction by means of spectacles or of a lens placed behind the sight-hole of the mirror. He should then seat himself opposite to and three or four feet away from the patient who is to gaze steadily at the darkened wall in front of him, looking towards the left side of the observer's head when the left eye is under examination and *vice versa* when the right is examined. By this means the optic disc is brought into the axis of vision of the observer. The latter must now throw the light from his mirror into the patient's eye, and, keeping the fundus illuminated, should move his head in various directions.

Indication 1. If he then see nothing more than the red reflex of the fundus or at most a *blurred* image of the disc, the eye is *Emmetropic* or *very slightly Myopic*. (*Vide* explanation, p. 25).

Indication 2. If he see the image of the disc and its vessels moving in the *same* direction as, or *with*, his own head, the patient is *Hypermetropic*. (*Vide* p. 28).

Indication 3. If the vessels in *one meridian only* are visible, and these move *with* the observer's head there is *Hyper-*

metropia in one meridian only, viz., in that at right angles to the one in which the vessels are visible, etc., = *Simple hypermetropic astigmatism*. (*Vide* explanation, p. 28 and compare with Chap. V.).

Indication 4. If the image of the disc and of the vessels appears to move in a direction *opposite* to, or *against*, that of the observer's head, the eye is *Myopic*. (*Vide* p. 28).

Indication 5. If the vessels are seen in *one meridian only* and appear to move *against* the direction of movement of the observer, there is *Myopia* in the meridian at right angles to that in which the vessels are visible = *Simple myopic astigmatism*. (*Vide* p. 28 and Chap. V.).

Indication 6. If the vessels are seen moving in one direction in one meridian, and in the opposite direction in the other meridian according to the accommodation of the observer, and his distance from the patient, there is *Mixed astigmatism*. This condition is, however, not easily detected by the direction of movement of the vessels.

Indication 7. If instead of the vessels of the disc moving evenly and regularly, they move slowly across the centre of the pupil, but rapidly and irregularly towards the periphery, giving the appearance of rotating bent spokes of a wheel, there is *Irregular astigmatism*.

EXPLANATIONS.

For many points in the following explanations, and especially in those relating to the change in size of the disc, Chap. X., I am indebted to my friend Dr. G. A. Berry.

Of all the rays, (*vide* Figs. 6 and 7), which diverge from any single point, *a* or *b*, of the fundus one only, *o'* or *o*,

passes without deflection through the centre of the crystalline lens. As many of the remainder as the size of the pupil will allow pass out in a cone of rays having a certain relation, according to the refraction, to the ray just mentioned. In order to get a view of any part of the fundus it is necessary that rays from *both* extremities of such portion should come to a focus on the observer's retina.




FIG. 6.

They thus leave between them an area x in which there are no rays from these two points. If the observer's eye be situated in this space it is evident that he cannot get an image of the *whole* disc. He will however receive *parallel* rays from some luminous *point* of the disc: or even cylinders from two very contiguous points

may run so closely together that on entering the observer's eye they might *theoretically* form an image of the space between them. *Practically*, however, such an image is rarely, if ever, obtained because the *absolute* suspension of accommodation necessary for its production is scarcely to be met with. We may therefore assume that for practical purposes no details of the fundus are visible with the mirror alone at some distance, and that therefore in *Emmetropia* the image of the disc is *blurred*.

In *Hypermetropia*, Fig. 6, the rays issuing from the extremities *a* and *b* of the disc emerge from the pupil as two cones (3 and 4) of rays which appear as if diverging from points *a, b*, situated behind the eye on prolongations backwards of the rays *o'* and *o*, respectively which from *a* and *b* pass through the centre of the lens. By the union of the rays thus prolonged back we get an *erect* image *a, b*, of the disc *a b*. (For the distance of this image behind the eye, *vide* p. 7, note to definition 6).

The cones 3 and 4 being formed of diverging rays, separate from each other only at a considerable distance from the eye. Some of the rays from each extremity of the disc, or at any rate those from two fairly distant portions of the same, will enter the observer's pupil, and, with the exercise of sufficient accommodation to overcome their divergence, will meet on his retina.

An object may be supposed as situated at the points whence appear to issue the rays entering the eye. The object therefore which the observer here appears to see is the *erect image* of the disc. Since the object thus apparently seen is *further* from the observer than the pupil, with

which it is, though perhaps unconsciously, compared, it seems to move with the observer's head, (*vide* explanation below).

In *Myopia*, Fig. 7, the rays issuing from the extremities, a b , of the disc emerge as two cones of rays which converge to points, a' b' , on their secondary axes, $O'O$, respectively, thus forming in front of the eye an *inverted* aerial image of the disc. If the observer be nearer the eye than where these rays unite, he will not get any image of the disc, since he receives only *convergent* rays which cannot come to a focus on his retina. If he, however, be far enough removed he will see the inverted image, a' b' , of the disc. The rays having converged to form this image, now cross and reach his eye, diverging (3 and 4) from the aerial image as if proceeding from an illuminated object placed at the same distance in front of the patient's eye.




FIG. 7.

The aerial image thus seen, being *nearer* the observer than the patient's pupil, moves against the observer's head. (*Vide* following explanation).

The apparent movement of the image of the disc is easily

explained when we remember that this image occupies a position further from the observer than the patient's pupil in H., and nearer to him than the pupil in M. This may be illustrated by holding two pencils vertically, one behind the other. If we suppose the *further* one to represent the image of the disc, and the *near* one the pupil, we have the respective positions of these objects in H. If, then, while moving the head laterally, we fix the eye on the *further* pencil, we find that this latter comes to be placed on the same side of the *near* one as that towards which we move. It *appears* therefore to travel *with* the observer.

If, however, we suppose the *near* one to represent the image of the disc and the *further* one the pupil, we have their respective positions in M. Then, on moving the head laterally, while regarding the *near* pencil, it will be seen that this latter comes to be placed to the *left* of the *further* one when we move to the *right* and *vice-versa*. It *appears* therefore to travel in a direction *opposite* to that of the observer.

The same explanation as that just given for the whole disc will apply also to vessels seen in different meridians.

In *Simple astigmatism*, we do not see vessels *at right angles* to the *emmetropic* meridian because rays from their transverse planes (Chap. V., p. 16) passing through the *emmetropic* meridian will (as was seen for simple *emmetropia*) not be visible. Whereas rays from the transverse planes of vessels *at right angles* to the *myopic* or *hypermetropic* meridian will, by passing through this meridian, produce in the former case an inverted, and in the latter an erect, image of the vessels. These follow the same rules of

movement as in simple myopia and hypermetropia respectively.

In *Compound myopic astigmatism*, if the observer use his accommodation, vessels at *right angles* to the *most* myopic meridian will be distinctly seen much *closer* to the patient's eye than will those in the opposite meridian. Rays from the transverse planes of the former converge and form an inverted image much sooner than do those from the latter, just as in high degrees of simple myopia.

In *Compound hypermetropic astigmatism* we find that for the vessels at *right angles* to the *most* hypermetropic meridian more accommodation is required at a given distance from the eye, than for those in the opposite meridian. A greater extent of the former than of the latter vessels is, however, at the same time visible. Rays from the former are more divergent and the cones take longer to separate, *vide* Fig. 6.

In *Mixed astigmatism* images of the vessels *may* be seen sometimes inverted and at other times erect according to the meridian through which the rays emerge, and varying with the observer's accommodation. The details are more visible than in emmetropia, but are nevertheless very *indefinite*.

CHAPTER IX.

RETINOSCOPY (KERATOSCOPY).

By means of what was, unfortunately, called "Keratotomy," we have a useful method of diagnosing errors of refraction with the ophthalmoscope mirror alone; and, what is perhaps more important, we can correct them with ordinary trial lenses quite independently of aid from the patient. This plan is particularly valuable where it is impossible to obtain, either suspension of the accommodation, or complete steadiness of the eye under examination, as well as the *certainly* of relaxation of the observer's own accommodation. Without these conditions, the estimation of refraction by the ophthalmoscope, as described in Chap. XI. is quite impracticable. As long ago as 1864 (*vide* "Anomalies of Refraction and Accommodation of the Eye," Donders, p. 490) Sir W. Bowman drew attention to "the discovery of regular astigmatism of the cornea, and the direction of the chief meridians, by using the mirror of the ophthalmoscope. . . . The area of the pupil then exhibits a somewhat linear shadow in some meridians rather than others."

Dr. Cuignet, of Lille, seems to have first systematised this method of examination, and, in 1874, published his conclusions in an article entitled "Keratotomy."

Dr. Parent, of Paris, published an article on this method of examination, in the *Recueil d'Ophtalmologie* for Feb. and

July, 1880, and Dr. Forbes has drawn attention to the subject in the *Oph. Hosp. Reports*, vol. x. For valuable information concerning the theory of keratoscopy I would refer the reader to Dr. Charnley's article in the *Oph. Hosp. Reports*, vol. x., part iii. Whilst for additional information respecting its mode of application, a reference to the *British Med. Journal* for Jan. 16, 1886, may prove useful.

The appearances, to be described, are due to the play of light and shade upon the *retina*, and are varied, in each case, by the refractive condition of the eye through which the rays of light reach, and issue from, the retina. Though the cornea, by means of its different curvatures, undoubtedly influences the appearances met with, yet, since it only acts as one of the refractive media, it was considered preferable to discontinue the use of the misleading word "Keratoscopy" and adopt, as suggested by Dr Parent, the term "Retinoscopy."

Rays of light from a distant lamp, falling on a *concave* mirror, issue from the latter convergingly, and crossing where they form an image of the lamp in front of the mirror, again diverge.

If, in front of a screen, we place a convex lens at such a distance that diverging rays from a concave mirror are brought to a focus exactly on the screen, there is formed the smallest and brightest possible image of the lamp, and the most sharply defined and densest surrounding shadow. If, then, we alter the position of the lens, it is found that the further it is removed from the point just mentioned, either towards, or away from, the screen, the *larger* becomes the area of light and the *feebler* is the illumination.

The increasing circles of diffusion render indistinct the line of demarcation between light and shade, and cause the latter to appear fainter. If, with the lens at different distances from the screen, the mirror be variously rotated, the area of light and shade will be seen, in *every* position of the lens, to move on the screen in a direction *opposite* to, or *against*, that in which the mirror is rotated. If we replace the screen and lens by the retina and dioptric system of the eye (p. 6, Def. 1) we have precisely similar results. The area of illumination on the *retina*, in *all* states of refraction, really moves in a direction *opposite* to that in which the mirror is rotated. But since this illuminated portion is seen through the transparent media of the eye the *apparent* direction of its movement will be influenced by the refraction of the eye under examination.

In the practice of Retinoscopy atropine certainly renders material assistance and without the aid of some mydriatic (atropine or homatropine) perfectly accurate results cannot be obtained. The observer should be seated opposite to, and 1 m. 20 cm. away from the patient. The lamp should be placed somewhat posterior to and above the patient's head. The eye which is not under examination should be carefully shaded by a screen, while the rest of the room should be rendered as dark as possible. The mirror used may be concave or plane. The concave mirrors in general use are of 22 cm. focal distance, and the following appearances are those met with in using such an instrument. If a plane mirror be employed the appearances will be the reverse of those here described. If the observer be not emmetropic, he must correct his own error of refraction.

The patient should look at the little hole in the mirror, while the light from it is thrown into his eye so that Retinoscopy is effected in his visual axis.

We then see, in the pupillary area, a bright image of the illuminated portion of the retina, bordered by a more or less linear shadow. If the mirror be now rotated around its various diameters, we shall see :—

1. That the direction of the *edge* of the shadow is at *right angles* to that meridian of the eye parallel to which the mirror is rotated.

2. That it travels along that meridian which is at *right angles* to the direction of its *edge*.

3. That it indicates the refractive condition of this last named meridian.

If, for example, the mirror be rotated around its vertical meridian from right to left or left to right, *i.e.* parallel to the horizontal meridian of the eye, the shadow will have its edge vertical; will travel in a horizontal direction across the area of the pupil; and will indicate the refraction of the horizontal meridian of the eye.

For any given meridian of the eye, the direction in which this shadow appears to move, *i.e.* whether with or against the mirror, depends upon, and furnishes the chief means of diagnosing the nature of, the refraction of that meridian.

If the rays issuing from the observed eye do not cross before reaching the observer, an erect image of the illuminated and shaded portion of the retina is obtained. This image appears to move in the direction actually taken by the illuminated area upon the retina, *viz.*, *opposite* to that in

which the mirror is rotated. This is the case in H., E., and *low* M.

If, however, the issuing rays cross before reaching the observer, and form, between him and the patient, an inverted *aerial image* of the illuminated and shaded portion of the fundus, then, since the illuminated area on the *retina* really moves *against* the mirror, the *aerial image* of the same will appear to move in the same direction as, or with, the mirror. This appearance is met with in cases of M. of 1 D. ($\frac{1}{40}$) and upwards: for if the observer, with good power of accommodation, be seated as stated, 1 m. 20 cm. from the patient, then, where the M. = 1 D. ($\frac{1}{40}$) the aerial image, being formed at 1 m. (40 ins.) from the patient, and 20 cm. (9 ins.) from the observer, will be easily perceived by the latter. Still more is this the case where, with increasing M., the image is formed nearer the patient. If, however, the M. be less than 1 D. ($\frac{1}{40}$), the convergent rays from the eye, either unite so close in front of the observer that no distinct image is received on his retina, or they meet behind him. In this latter case, the rays *not having crossed* before reaching the observer's eye, the image of the shadow is seen to move *against* the mirror. The way in which this condition is distinguished from H. or E., is explained further on (p. 37). If the eye under examination be to such an extent myopic that its "far point" is situated at the position of the image of the lamp, say 25 cm. in front of the mirror, or 95 cm. from the eye, then the diverging rays from this image will come to a focus exactly on the patient's retina (Def. 5, p. 7) and there form the smallest and brightest possible image of the lamp.

The further the departure from this degree, which we may call 1 D. ($\frac{1}{40}$) of M., the larger and feebler, as was seen on the screen, will be the illuminated portion of the retina.

The difference in the degree of luminosity of this reflex can be easily appreciated, and furnishes a means whereby the amount of error may be approximately estimated: so that, speaking generally, we may say, the feebler the illumination the higher is the ametropia. An attempt has been made to use, for the same purpose, the difference in intensity of the shadow, but the appreciation of this is difficult, and it therefore does not afford an altogether reliable source of information.

If, from some distance, say 1 m. 20 cm., we examine an eye with the mirror alone, we find that the higher the H. or M., the smaller is the image which we obtain of the disc: so that in very high degrees, we see not only the whole disc, but also some of the surrounding fundus in the pupillary area. In E., on the other hand, so large is this image that only a small portion of it is visible at one time. With equally rapid rotations of the mirror then, the light would have to travel much faster over the large image of the latter, than the small image of the former, condition. This difference in the *rate* of movement in the various states of refraction, was pointed out by my friend Dr. Charnley: it constitutes, especially when taken together with the *luminosity* of the image, probably the best of the means at our disposal for estimating, by this method without glasses different *degrees* of ametropia.

In E. and the lower degrees of H. and M., so little is

seen of the large image of the illuminated area and the surrounding shadow, that the small portion of the latter, visible at any one time in the pupillary area, appears approximately *linear*; while, although with the increasing degrees of H. and M., the nearly circular area of illumination on the retina also enlarges, yet so much diminished is the image of the same, that more of the circumference of the surrounding shadow is visible at one time in the area of the pupil; it appears, therefore, more *crescentic*, while it becomes at the same time narrower.

Taking then the direction of the movement as denoting the *kind* of error; and the rate of movement, degree of luminosity, and curvature of the shadow, as indicating approximately the *amount* of the same, we may summarise as follows:—

1. If the image of the shadow appear to move in the direction in which the mirror is moved; and, if the rapidity of movement and curvature of the shadow are the same in all meridians, we have a case of *simple M.*
2. If the image of the shadow appear to move in the opposite direction to that in which the mirror is moved: and, if the rapidity of movement and curvature of the shadow are the same for all meridians, we have to do with either H., E., or *low M.*
3. The *slower* the movements of the image, the *feebler* the illumination, and the more *crescentic* and narrower the shadow, the *higher* is the H. or M.
4. A *difference*, in two opposite meridians, either of the direction or rapidity of movement, or of the curvature, of the shadow, indicates *Astigmatism*. These two dissimilar

shadows, moving at right angles to each other, either one vertically and the other horizontally, or both obliquely, indicate the position of meridians of greatest and least refraction.

The situation of the principal meridians can be ascertained by noting whether the movement of the shadow in one meridian (particularly when nearly over-corrected) is complicated with movement in another; when there is no such complication it is probable that in most cases retinoscopy is being practised in one of the principal meridians. This movement of the opposite shadow also takes place in eyes characterised by irregular astigmatism.

It must be carefully noted that the direction of movement of the *shadow*, as compared with the *mirror*, is the *reverse* of that of the *disc* or vessels as compared with the *observer's head* (*vide* Chap. VIII.).

The *amount* of error may easily be measured by ordinary trial lenses. A little practice soon enables one to make an approximate estimate of the refraction, and to start with a lens nearly correcting the ametropia. At first, however, it is necessary to proceed carefully by quarter dioptries from zero. It will be well to remember (page 35) that in high degrees of ametropia the reflex is often so feeble that there is great difficulty in determining the direction of the shadow. This is perhaps best ascertained by putting up successively a strong convex and a strong concave lens (4 D. or 6 D.). One of these will probably so far correct the ametropia that the reflex becomes bright and the shadow discernible. The further correction is then easily accomplished. Again in M. below 1 D. and

in E., though the reflex is bright, there is no definite shadow movement. This condition is easily distinguished from the preceding where the reflex is dull.

Suppose now that we have first to deal with a case where the shadow moves *against* the mirror, or where, with a bright reflex, there is no perceptible movement of the shadow. We then know that the eye has either H., E., or weak M. But as we cannot tell which is present, we place in front of the eye *convex* lenses increasing in strength till one is reached which causes the shadow to move *with* the mirror. As soon as this occurs we know that M. of 1 D. is present. If this result is obtained with the weakest glass used (+ 0.25 D.), then this latter will account for 0.25 D. of the M., while the remaining 0.75 D. of M. must have already existed in the eye. In the same manner, if the weakest glass required to reverse the shadow were either + 0.5 D., + 0.75 D., or + 1 D., there must have previously existed in each of these cases respectively, 0.5 D. of M., 0.25 D. of M. or E.

On the other hand, if with + 1 D. the shadow still moves *against* the mirror it shows that the eye is hypermetropic. If the weakest glass which reverses the movement is + 1.25 D., it shews that the H. is 0.25 D. Had the eye been emmetropic the shadow would have been reversed by + 1 D., while if H. to the extent of 0.5 D. had been present, then the + 1.25 D. would not have over-corrected it by 1 D., and the shadow would not have been reversed. Similarly with the higher lenses. If + 4.5 D. were the weakest which reversed the shadow, it would shew that artificial M. of 1 D. had been produced, and the H. over-

corrected by 1 D. The H. therefore really present would be 3.5 D.

In the class of cases just illustrated then, where the original shadow moves *against* the mirror, the actual refraction is indicated by deducting + 1 D. from the weakest *convex* lens which causes the shadow to move *with* the mirror.

In the second place suppose we have to deal with a case where the shadow moves *with* the mirror. From this fact we know that there must be present M. of at least 1 D. (p. 34). We then begin with the weakest *concave* lens and proceed by quarter dioptres till one is reached with which the shadow *ceases* to move with the mirror. As soon as this is the case, say with the weakest lens we try, viz., - 0.25 D., it shews that there cannot now be 1 D. of M. left uncorrected. The original M. must therefore have been exactly 1 D. Had it been even as much as 1.25 D., then the - 0.25 D. would have left 1 D. of M. uncorrected, and the shadow would have continued to move with the mirror. In the same way, if - 2.25 D. were the weakest glass with which the shadow ceased to move with the mirror, it would shew that the original M. actually present was 3 D. This is evident, because with - 2 D. the shadow was still with the mirror, and therefore, 1 D. of M. was left uncorrected, while with - 2.25 D. the shadow ceased to move with the mirror, and therefore, there could not have been 1 D. uncorrected. In other words the M. could not have been as much as 3.25 D.

We see, therefore, that in this latter class of cases where the original shadow moves *with* the mirror, the actual

refraction is indicated by deducting $+ 1$ D. from the strongest *concave* lens with which the shadow continues to move *with* the mirror.

This deduction of $+ 1$ D. from a lens, whether it be convex or concave, has the same result as adding $- 1$ D. to it. The signs plus (+) and minus (−) as applied to the trial lenses are algebraic terms, and the addition or subtraction of them follow the ordinary rules of algebra. For the sake of uniformity and simplicity it is better, as suggested by Dr. Casey Wood in the *Journal of Ophthalmology, Otology, and Laryngology* for 1889, that the same rule shall be made to apply to both convex and concave glasses.

If the same lens produces similar results for all meridians there is no astigmatism. But if, when the movement is reversed in one principal meridian it retains its original direction in the other, this proves astigmatism to be present. Each of the two principal meridians must then be corrected separately. This may be effected by correcting the refraction in one principal meridian by a spherical glass, and then, leaving it in position, correcting it in the other by means of an additional cylinder at right angles to this latter meridian. A more convenient method, however, and one generally adopted, is to correct the refraction in each of the principal meridians by means of separate spherical lenses. The degree of astigmatism is indicated by the cylinder in the former method, and by the difference between the two sphericals in the latter plan.

Having, by the above method, measured the refraction, the V. may be tested by the glasses thus indicated. These

will give the best results in the majority of cases. Some patients, however, see better with lenses rather weaker, others with them stronger than the retinoscopy results indicate. It will, therefore, be occasionally necessary to make some modification in the glasses thus found, and especially is this likely to be the case in myopia.

If the refraction is the same in all meridians, a spherical lens will suffice.

With only one meridian ametropic, a cylindrical lens, with its axis at right angles to this meridian, will be necessary.

When there is either H. or M. in both of the principal meridians, the least ametropic meridian is corrected by a spherical lens. An additional cylinder, of the same sign, and equal to the difference in refraction, between the two meridians, is then placed at right angles to the opposite meridian.

Where there is H. in one principal meridian and M. in the other, the former is corrected by a *convex* spherical lens. An additional *concave* cylinder, equal to the difference in refraction between the two meridians is then placed at right angles to the myopic meridian.

The glasses which give the best results while the patient is under atropine, sometimes also give the best V. when the effect of that drug has passed off. In other cases again, some modification may be necessary. A convex lens not infrequently requires reducing in strength. A concave lens seldom requires any alteration. With spherico-cylinders any modification necessary may almost invariably be effected by means of the spherical lens alone.

If the patient cannot be seen after atropine has passed

off, allowance may be made for its action (*vide* Chap. XIII.) by deducting a certain convex lens (varying according to the amount which it is deemed advisable to allow for atropine) from the *spherical* glass of the spectacles with which best V. is obtained. If a preliminary trial with glasses is not possible, the deduction may be made from the figures shewing the actual refraction. The same amount must be deducted from each meridian so that the relative refraction may not be altered.

When the glasses are to be worn constantly, it is a good plan to let the patient continue atropine once daily until he has begun to wear them.

In conclusion it may be well to recapitulate some of the more important points.

(i.) The observer must always be seated at the same distance from the patient, viz., about 1 m. 20 cm.

(ii.) The mirror may be concave or plane, but the indications are of course exactly opposite with the two mirrors.

(iii.) The accommodation and the iris being paralysed by homatropine or atropine, a screen should be placed over one eye and the patient should look at the sight-hole in the mirror.

(iv.) The situation of the principal meridians should then be roughly ascertained, and the degree of refraction tested with glasses. It is more convenient to test each meridian separately and with spherical glasses only in the first instance.

(v.) If the shadow move against the mirror, or if with a *bright* reflex there is no definite shadow movement, we have a case of E., H., or *weak* M.

(vi.) If the shadow appear to move with the mirror, the eye is myopic.

(vii.) To ascertain the actual refraction while the eye is still under atropine :—

Deduct $+ 1$ D.

(a) From the weakest *convex* glass which *reverses* the shadow if it originally moved against the mirror.

(b) From the strongest *concave* glass with which the shadow *continues* to move *with* the mirror if it originally had this direction.

(viii.) The *feebler* the luminous reflex and the *slower* the movement of the shadow across the pupillary area, the greater is the amount of ametropia.

To illustrate the method of working retinoscopy, a few examples of astigmatism are appended. The cross lines, which for the sake of simplicity are placed vertical and horizontal, represent the two meridians. The figures refer to the meridian at the end of which they are placed. Those in the first brackets indicate the retinoscopy results as measured by trial lenses. The figures in the second brackets denote the actual refraction, and are derived by deducting $+ 1$ D. from the previous numbers. While those in the third brackets shew a further deduction of $+ 1$ D. for the action of atropine. The latter deduction does not at all imply that this is the amount which should always, or even generally, be deducted, but is merely given by way of illustration.

The glasses in the first brackets correspond to the actual refraction under atropine as shewn by the figures in the above mentioned second brackets. The glasses in the

second brackets shew the deduction of $+ 1$ D. for atropine, and correspond to the figures in the above mentioned third brackets. This deduction may also be made from the *spherical* lens of the previous glasses or of those with which best V. was obtained under atropine.

To prevent confusion the following examples are given only in dioptric measurements.

The sign \subset means "combined with."

"G" = glasses.

CASE I.—Simple hypermetropic astigmatism.

$$\begin{array}{|l} (+2) (+1) (0) \\ \hline (+5) (+4) (+3) \end{array} = G. (+1 \text{ s. } \subset +3 \text{ c. vert.}) (+3 \text{ c. vert.}).$$

CASE II.—Compound hypermetropic astigmatism.

$$\begin{array}{|l} (+3) (+2) (+1) \\ \hline (+5) (+4) (+3) \end{array} = G. (+2 \text{ s. } \subset +2 \text{ c. vert.}) (+1 \text{ s. } \subset +2 \text{ c. vert.}).$$

CASE III.—Simple myopic astigmatism.

$$\begin{array}{|l} (-3) (-4) (-5) \\ \hline (+1) (0) (-1) \end{array} = G. (-4 \text{ c. hor.}) (-1 \text{ s. } \subset -4 \text{ c. hor.}).$$

CASE IV.—Compound myopic astigmatism.

$$\begin{array}{|l} (-3) (-4) (-5) \\ \hline (+0.5) (-0.5) (-1.5) \end{array} = G. (-0.5 \text{ s. } \subset -3.5 \text{ c. hor.}) (-1.5 \text{ s. } \subset -3.5 \text{ c. hor.}).$$

In Cases III. and IV., it is most probable that the glasses found under atropine to be the best, as indicated in the first brackets, are those which would be ordered for permanent use without making any allowance for atropine.

CASE V.—Mixed astigmatism.

$$\begin{array}{c} | (+0.75) (-0.25) (-1.25) \\ \hline (+3) (+2) (+1) \end{array} = G. (+2 \text{ s. } \ominus -2.25 \text{ c. hor.}) (+1 \text{ s. } \ominus -2.25 \text{ c. hor.}).$$

CASE VI.—Mixed astigmatism.

$$\begin{array}{c} | (+1.5) (+0.5) (-0.5) \\ \hline (+4.5) (+3.5) (+2.5) \end{array} = G. (+0.5 \text{ s. } \ominus +3 \text{ c. vert.}) (+2.5 \text{ s. } \ominus -3 \text{ c. hor.}).$$

Suppose that in this case the best V. was obtained, not by the glasses indicated in the first bracket, but by + 3 D. *c. vert.* This would shew that, under atropine, the vertical meridian was emmetropic and the horizontal meridian, hypermetropic 3 D., thus:—

$$\begin{array}{c} | 0 (-1) \\ \hline +3 (+2). \end{array}$$

After deducting + 1 D., therefore, for atropine from each meridian, as shewn in the brackets, the final glasses would be + 2 s. \ominus - 3 c. *hor.* The same result would be obtained by placing - 1 D. in front of the + 3 D. *c.*, thus:—
- 1 D. s. \ominus + 3 D. *c. vert.*, but it is better to adhere to the rule of making the *spherical* glass *convex*.

CHAPTER X.

INDICATIONS AFFORDED BY MIRROR AND OBJECT LENS.
"INDIRECT METHOD."

THIS method of diagnosing Hypermetropia and Myopia was first pointed out by Mr. Hutchinson (*Ophth. Hosp. Rep.*, vol. iv., p. 189) and Mr. Couper (*Med. Times and Gaz.*, Jan. 30, 1869) has shown how Astigmatism may also be recognised by it. It is desirable that atropine should have been applied, for otherwise an alteration in the pupil may easily lead to the supposition that the disc has changed in size.

In this mode of examination the observer must place the object lens *as close as possible* to the patient's eye; then, keeping the optic disc steadily in view, he must gradually withdraw the lens to the distance of a few inches: when—

Indication 1. If the disc remain the *same size* throughout, the patient is *Emmetropic* (*vide* expl. p. 47).

Indication 2. If the disc *diminish* in size on withdrawing the lens the patient is *Hypermetropic*, and the amount is proportional to the rapidity of diminution (*vide* expl. p. 48).

Indication 3. If the disc *decrease* in size *in one meridian only*, the patient is *Hypermetropic* in *that meridian only* = Simple hypermetropic astigmatism.

Indication 4. If the disc diminish in all meridians, but *more* in *one meridian* than the others, there is *Hypermetropia*

in all meridians, but the greatest is in that meridian which decreases most = Compound hypermetropic astigmatism.

Indication 5. If the disc *increase* in size on withdrawing the lens there is *Myopia*, and the amount is proportional to the rapidity of increase (*vide* expl. p. 49).

Indication 6. If the disc *increase* in *one* meridian only, there is *Myopia* in that meridian alone = Simple myopic astigmatism.


Indication 7. If the disc *increase* in every direction, but *more* in *one* meridian than another, there is *Myopia* in all meridians but the greatest is in that meridian which increases most = Compound myopic astigmatism.

Indication 8. If the disc *increase* in *one* meridian, and *diminishes* in the *opposite* one, there is *Myopia* in the former, and *Hypermetropia* in the latter = Mixed astigmatism.

Explanation of change in size of the disc.—These changes in size of the image of the disc will be evident on remembering that *the relative sizes of image and object are as their distances from the lens*. To find the distance of the image from the lens, we have the formula $\frac{1}{b} = \frac{1}{f} - \frac{1}{a}$, where b = the distance of the image from the lens, f , = the focal length of the lens, and a = the distance of the object. In each of the following examples the focal length of the lens is 4 cm.

In *Emmetropia* the rays issuing from the disc emerge from the eye *parallel* as if proceeding from an object situated at an infinite distance (Def. 3, p. 6). It is of this supposed object that we get an image by means of a convex lens held in front of the patient's eye. It matters not, therefore, where

this lens is held, the *parallel* rays from the object will always unite to form an image on the opposite side of the lens at its principal focus (Chap. I., *b*, p. 3). The relative distances of image and object from the lens remaining *constant*, the size of the image of the disc in Emmetropia does not vary with movements of the lens.



In *Hypermetropia*, Fig. 8, * rays issuing from the disc emerge from the eye as if proceeding from an object, *b a*, situated at a certain distance behind it (Def. 6, p. 7).

It is of this supposed object, *b a*, that we gain an *image*, *a, b*, or *I'*, by means of a lens held in front of the patient's eye.

Suppose this lens, *L*, to be at 6 cm. from the object, we find by the formula that the distance of the image *a, b*, is 12 cm. ($\frac{1}{b} = \frac{1}{4} - \frac{1}{6} = \frac{1}{12}$ or $b = 12$.)

If we now withdraw the lens from the object (and also from the eye) till it is 12 cm. from the former as at *L'*, the distance of the image, *I'*, (between the dotted lines) will then be 6 cm. ($\frac{1}{b} = \frac{1}{4} - \frac{1}{12} = \frac{1}{6}$ or $b = 6$.)

FIG. 8.

The ratio of the distance of the image from the lens as compared with that of the object from the lens being *greater* in the first case than in the second, so is the size of the image. In *Hyper-*

* In figs. 8 and 9, the distances of the lenses from the objects and images are drawn to scale $\frac{1}{2}$ cm. = 1 cm.

metropia on withdrawing the lens from the eye, the image of the disc *diminishes* in size.

In *Myopia*, Fig. 9, rays from the disc emerge from the eye convergently, and form, at a certain distance in front of it, an inverted aerial image, $a\ b$, of the disc. This latter we have now to regard as the object, o , whose image, I or I' , we obtain by means of a convex lens, L or L' , interposed between it and the patient's eye, E . A lens, thus placed, intercepts the rays, Bb , Aa , (which, to avoid confusion, are the only ones shown of all those) which converge towards the points b and a respectively of this object, o . The rays being thus rendered more convergent, produce an image I or I' , whose extremities will be bounded by the lines passing from either end respectively of the object, o , through the centre of the lens, L or L' .




FIG. 9.

In this case, object and image are on the *same* side of the lens. The object here, being on the opposite side of the lens from the direction in which the rays proceed, it is customary, in order that the same formula may hold good in all cases, to regard $\frac{I}{a}$ as a *negative* quantity. For the distance of the image from the lens then, we have the formula $\frac{I}{b} = \frac{I}{f} - \left(-\frac{I}{a} \right) = \frac{I}{f} + \frac{I}{a}$.

If we first place a lens, L , 12 cm. nearer the eye than where the object o would be formed, we have the image I

at 3 cm. from the lens ($\frac{1}{b} = \frac{1}{4} + \frac{1}{12} = \frac{1}{3}$ or $b = 3$). If we now withdraw the lens *from* the eye, *i.e.*, towards the object till it is within 6 cm. of the latter, as at L' , the distance of the image I' will be nearly 2 cm. ($\frac{1}{b} = \frac{1}{4} + \frac{1}{6} = \frac{5}{12}$ or $b =$ nearly 2). But as the ratio of the distance of the image from the lens, as compared with that of the object from the lens, is greater in the second than the first of these examples, so is the size of the image. In Myopia, therefore, on approaching the lens to the eye the image diminishes in size, and on *withdrawing* it from the eye the image *increases*.

The above explanation holds good for *Myopia* only so long as the lens is not withdrawn beyond the "far point" of the eye *plus* its own focal distance: for *Hypermetropia* only so long as the focal power of the lens is greater than the degree of Hypermetropia. The exception to this latter condition does not occur in practice if a lens of 3 ins. focus be used and need not therefore be considered. In the case of very high degrees of Myopia, however, if the lens be further from the eye than the aerial inverted image of the disc *plus* its own focal distance, an erect image of the disc is formed between the lens and the observer, the variations in size of which are subject to the same rules as those described for the inverted image in Hypermetropia.

CHAPTER XI.

ESTIMATION OF THE REFRACTION BY MEANS OF LENSES IN
THE OPHTHALMOSCOPE.

IN estimating refraction by this method the patient and observer must be seated side by side facing in opposite directions. Their heads and the lamp should be on the same level; the latter being placed on the same side of the patient as that under examination, a little out from, and slightly behind the position of, the patient's ear. If now the heads be inclined laterally towards each other, the eyes of the corresponding sides will come opposite one another, while the noses and mouths will be left free for breathing. By this means the sight-hole of the mirror may generally be placed in the position which, when ordered, the glasses will occupy; the distance of these from the eye need therefore not be taken into consideration when prescribing from measurements thus obtained. Should it however be undesirable to approach thus close, the distance from the patient must be taken into account, and the glasses ordered will have to be somewhat *weaker* in *myopia* and *stronger* in *hypermetropia* than the lens necessary to see the fundus would indicate. In estimating refraction by the ophthalmoscope it is absolutely essential that the accommodation of both patient and observer be *completely* suspended. In order to secure the former we must place the

patient opposite a dark wall or curtain, or employ atropine. For the observer voluntarily to relax his own accommodation while looking at an object so near to him as the patient's eye, requires much practice. The best way of overcoming the difficulty is to regard the fundus as situated some hundreds of yards distant. This habit may be acquired by looking alternately at a spot on the window pane and then at some remote object: or by gazing vacantly at the page of a book until the type disappears; then, having noticed the sensation produced on thus relaxing the accommodation, endeavour to induce the same condition while observing a fundus.

In the following indications it is of course necessary that the observer be fully cognisant of the state of his own refraction, and in the following cases he is supposed to be emmetropic, (if not, *vide* note, p. 54). With accommodation, then, entirely suspended, the eye of the emmetropic observer is adjusted for *parallel* rays.

Indication 1. If therefore he can see distinctly the details of any fundus, the rays issuing thence must be *parallel* and the eye *emmetropic*.

NOTE. Though this appears to contradict what has been said concerning seeing the disc in emmetropia (Chap. VIII., p. 25), yet we must remember that when *close* up to the patient's pupil, the observer *does* receive rays from the two extremities of the disc. Moreover if the observer endeavour to accommodate for the patient's eye, he will find it impossible to do so, since, if he is as close as he should be, the latter, as explained by Mr. Power, is nearer to him than his "near point." The effort is therefore not main-

tained, and as when the accommodation is partially suspended the disc begins to appear, the former is unconsciously *altogether* relaxed and a distinct view is obtained.

Indication 2. If, however, the details of the fundus cannot be seen except by the intervention of a *concave* lens, this proves the patient to be *Myopic*. The converging rays from such an eye (Def. 5, p. 7) require a concave lens to render them parallel for perception by the emmetrope. The strength of the lens which effects this change of direction indicates the degree of M., for it is evident that the same lens which renders parallel the converging rays issuing from the retina, will bring parallel rays to a focus on the retina of the same eye.

Indication 3. If again, with suspended accommodation, the details of the fundus are not visible without the aid of a *convex* lens, this indicates the patient to have *Hypermetropia*. The diverging rays from a hypermetropic eye require a convex lens to render them parallel. The strength of the lens necessary for this purpose denotes the amount of H., for the lens which renders parallel, rays proceeding from a retina, will also bring parallel rays to a focus on the same retina.'

Indication 4. If the vessels in one meridian are seen without any lens while a convex or concave lens is required for those in the other meridian, it is a case of *Simple astigmatism*. The emmetropic meridian is the one at *right angles* to the vessels seen without any lens (Chap. V., p. 16).

Indication 5. If for the two opposite meridians, we require either two convex or two concave lenses of *unequal* power, we have to deal with a case of *Compound* astigma-

tism: hypermetropic or myopic respectively. The greatest error of refraction in each case is at *right angles* to the vessels for which the strongest lens is necessary.

Indication 6. If for vessels in one meridian we require a *convex* lens, while for those in the other a *concave* is necessary, it indicates *Mixed* astigmatism. The hypermetropic meridian is at *right angles* to the vessels seen best with a convex lens: and the myopic to that for which a concave lens is necessary.

In all these forms of astigmatism then, we see that the lens which, with *absolutely* suspended accommodation, gives the best view of vessels in *one* meridian, measures the amount of error which exists in the *opposite* meridian.

NOTE. Should the observer not be emmetropic, yet, if he correct his own error with spectacles, he is in the same position as an emmetropic observer. If he, however, prefer to estimate refraction without his glasses he must make an allowance for his defect by means of the lenses behind the sight-hole of the mirror. To see the details of an *emmetropic* fundus, a hypermetropic or myopic observer *starts* with a lens corresponding to the amount of his own error. For any particular fundus, therefore, the glass required by him must be, to the extent of his H. or M., *more* convex or concave respectively than would be necessary for an emmetrope to see the same fundus. Consequently the *patient* will always have that amount *less* H., and *more* M.; or *more* H., and *less* M., respectively than the same lens would indicate if the observer were emmetropic; we have then the

RULE II. That the observer must *deduct* the amount of his own H. or M. from the lens which enables him to see distinctly any particular fundus.

TABULAR VIEW OF THE STATE OF REFRACTION AS INDICATED BY THE VARIOUS METHODS.

TEST TYPES.				OPHTHALMOSCOPE.		
DISTANT VISION.	NEAR VISION.	RAYS IN FAN.	MIRROR ALONE.	INDIRECT METHOD.	Lens necessary in Ophthalmoscope without accom.	REFRACTION.
$\frac{6}{6}$ ($\frac{20}{20}$) By accom. may be $\frac{6}{6}$ but is often less.	Sn. 1, 5 in. to 20 in.	All clear.	Movement of <i>image</i> of O, D, and vessels compared with <i>Head</i> . Red reflex only.	Movement of <i>Image</i> of O, D. on <i>withdrawing</i> lens. Constant in Size.	None.	E.
	Sn. 1, or larger, but not close up.	All clear by accom.	Erect; moves with.	Against; same rate in all meridians.	Convex.	H.
Not $\frac{6}{6}$	Sn. 1, (or 2) but not so far off as distance marked.	None.	Inverted; moves against.	With; (except low M.) same rate in all meridians.	Concave.	M.
Not $\frac{6}{6}$	Larger series (or smaller with difficulty).	One clear at rt. ang. to Em. meridian.	Vess. parallel to Em. merid. move with.	In two opp. merid. Both against at diff. rates.	Vess. Par. to Em. merid. = + Vess. Par. to Hc. merid. = O.	H. As.
Not $\frac{6}{6}$			Vess. parallel to Em. merid. move against.	In two opp. merid. one against, one with.	Vess. Par. to Em. merid. = - Vess. Par. to Mc. merid. = O.	M. As.
		Varying with accom.	Vess. at diff. distances, all move with.	In two opp. merid. Both against at diff. rates.	Decreases all round, but more so in one meridian.	C.H. As.
		None.	Vess. at diff. distances, all move against.	In two opp. merid. Both with at diff. rates.	Increases all round but more so in one meridian.	C.M. As.
Variable.	Larger only.	One at rt. ang. to Hc. merid. by accom.	Some vess. with others against.	In two opp. merid. one against, one with.	One merid. + Opp. merid. -	Md. As.

Abbreviations:—E. Emmetropia; M. Myopia; H. Hypermetropia.
H. As. Simple Hypermetropic Astigmatism; C. H. As. Compound Hypermetropic Astigmatism.
M. As. Simple Myopic
Md. As. Mixed Astigmatism; Vess. Vessels; Par. Parallel; Diff. Different; O. D. Optic Disc.

CHAPTER XII.

TEST GLASSES AND TYPES AS APPLIED TO THE ESTIMATION
OF REFRACTION.

WHEN ascertaining with the distance types the state of the patient's refraction, we must be absolutely certain that his accommodation is completely relaxed. In myopes it generally is so, but in hypermetropic and astigmatic patients it is frequently a troublesome and misleading factor. If, from the varying and inconsistent statements of the patient, it is suspected that he does not relax his accommodation, he may do so more easily if he close his eyes between each change of glasses, not opening them until the fresh ones are *in situ*. Another method of assisting the patient is to let him put on a pair of + 4 D. glasses for ten or fifteen minutes, and then, without removing, gradually neutralise them by stronger and stronger concave lenses placed in front, until those are found with which the patient can see in the distance as well as without any glass. The *difference* then, between the concave lens thus required and + 4 D. ($\frac{1}{10}$), gives the manifest hypermetropia.

If, however, there is any uncertainty as to the accommodation being relaxed, it is better to have recourse to atropine. In all cases each eye must be tested separately.

One caution must be especially borne in mind, viz.: *never*, in any doubtful case, to commence testing vision with *con-*

cave glasses, for to see with such lenses it is necessary for all, except those whose myopia is equal to or less than the glasses being used, to employ their accommodation, and when once called into activity, this is not easily suspended.

Let the patient be placed at 6 m. (20 ft.) from the types. We have already seen what, with test types, are the indications of emmetropia and simple presbyopia (p. 21).

HYPERMETROPIA.

If the patient read $\frac{6}{6}$ ($\frac{20}{20}$) perfectly without glasses, still *this fact does not exclude the presence of Hypermetropia*, the presence of which is proved if he can read the distance types as well with, as without, *convex* lenses. The *strongest* glasses which are thus tolerated indicate the degree of *manifest* hypermetropia (Hm.) and they *at least* must be ordered for close work. In young adults, and in the higher degrees of H., we must for this purpose even give glasses 1 D. or 1.50 D. ($\frac{1}{40}$ or $\frac{1}{24}$) stronger. In order to give rest to the ciliary muscle, it is advisable for constant use to prescribe at any rate the glasses which correct the *manifest* hypermetropia. If the H. be measured under atropine, the glasses ordered must be 1 D. (or possibly even 1.50 D.) *weaker* than the total amount thus found to exist (*vide* Chap. XIII.). These should be suitable for all purposes, though for distant V. weaker ones are sometimes necessary.

The amount of hypermetropia which is apparent only after atropine is applied, is called the latent hypermetro-

pia (Hl.) and added to the Hm. gives the total hypermetropia (Ht.). The Hm. steadily increases and the Hl. steadily decreases as age advances till at last the Hm. = Ht.

MYOPIA.

As we have already stated (Ind. 3, p. 21), if the patient can read the *finest* type to within 4 ins. or 5 ins. of his eye while at the same time his distant V. does not exceed $\frac{6}{24}$ ($\frac{20}{70}$) and is probably not $\frac{6}{60}$ ($\frac{20}{60}$), he is *myopic*. An indication of the degree may be obtained by observing the furthest point at which either of the smallest types can be read. Provided it be nearer than that for which it is marked, the distance of this point gives the focal length of the lens required to neutralise the M.

e.g.—If Sn. 1 be legible only as far off as 12 cm. (5 ins.) the M. is measured by the concave lens having this focal length, viz., 8 D. ($\frac{1}{8}$). Or again, if No. 4 Sn. marked for 1 D. (40 ins.), can be read only at 50 cm. (20 ins.), the M. = 2 D. ($\frac{1}{20}$).

In testing the distant V. with glasses we commence with the *weaker* concave glasses, and work up to the stronger. In doing this, we cannot be too careful in ascertaining which is the *weakest* glass that neutralises the M. and gives the best V., whether this be $\frac{6}{6}$ ($\frac{20}{20}$) or less. The glass thus found gives the measure of the M. and may generally be ordered for distant V. Such glasses may also be given for close work when, *with good accommodation*, the M. does not exceed about 6 D. or 8 D. ($\frac{1}{6}$ or $\frac{1}{8}$). In most cases where the M. is higher, and in all where the accommoda-

tion is feeble, we must order *weaker* glasses for *close* work. The strength of these may then be found in the following manner, viz: From the neutralising lens *deduct* the strength of the glass whose focal length equals the distance at which we wish the patient to work.

e.g.—With M. = 10 D. ($\frac{1}{4}$) we wish the patient to do work at 40 cm. (16 ins.). From 10 D. ($\frac{1}{4}$) we deduct therefore the lens whose focal length is 40 cm. (16 ins.), viz: 2.50 D. ($\frac{1}{16}$), and the glasses ordered will be 10 D.—2.50 D. = — 7.50 D. ($\frac{1}{4} - \frac{1}{16} = \frac{1}{6\frac{2}{3}}$).

ASTIGMATISM.

If the V. cannot be brought up to $\frac{6}{8}$ ($\frac{20}{20}$) with any *spherical* glasses, we probably have to do with a case of *Astigmatism*. In dealing with this error, as with all others, each eye should be tested separately.

If, from the previous examination with the ophthalmoscope, we have, diagnosed,—

1. *Simple* astigmatism, we can proceed to test with the fan of rays. If our surmise has been correct, the patient should now see quite distinctly, only the line at *right angles* to his emmetropic meridian (Chap. V., p. 16). If this line were vertical, it would therefore denote that the meridian *parallel* to it was either hypermetropic or myopic; we should then ascertain what cylinder, with its axis at *right angles* to this latter meridian is required to correct it: *i.e.*, to render clear the horizontal line. If the correction thus found gives V. = $\frac{6}{8}$ ($\frac{20}{20}$) it may be ordered for constant use;

but if it do not, then, as where none of the rays are distinct, and more particularly if first one then another is most clearly seen, we must not hesitate to employ atropine.

It matters not now, whether we have to deal with—

2. Compound or mixed astigmatism. We have merely to ascertain what *spherical* lens clears *one* of the rays, and then, leaving this glass *in situ*, try what cylinder, with its axis at *right angles* to the line thus cleared, renders equally distinct the ray in the opposite meridian. This spherico-cylindrical correction, *after due allowance for atropine* (Chap. XIII.), is ordered for constant use.

3. Another less scientific, though practically useful plan, is to substitute the test-types for the lines, and having found the spherical lens which gives the best V., try what additional cylinder is required.

4. If each meridian has been measured separately with spherical glasses, either for the fan, or in the ophthalmoscope for the fundus, we shall have to calculate what spherico-cylinder is required. (*a.*) For *compound* astigmatism we generally give the spherical lens, which corrects the meridian of least error, and then add the cylinder (concave for myopic, and convex for hypermetropic astigmatism), whose strength equals the difference between the two meridians. (*b.*) In *mixed* astigmatism the difference between the sphericals also gives the degree of astigmatism and the strength of the cylinder required. It is advisable in ordering this combination to prescribe the spherical lens convex and the cylindrical lens concave.

e.g.—With vertical meridian myopic 2 D. ($\frac{1}{20}$) and the

horizontal meridian hypermetropic to the same extent, the *difference* between them = 4 D. ($\frac{1}{10}$). If, therefore, we give *plus* 2 D. sph. ($\frac{1}{20}$), we must add *minus* 4 D. cyl. ($\frac{1}{10}$) with its axis *horizontal* (i.e. at *rt. angles* to the myopic meridian, Chap. I., p. 4). But if we gave *minus* 2 D. sph. ($\frac{1}{20}$), we should require in addition *plus* 4 D. cyl. ($\frac{1}{10}$) axis *vertical*. The action of atropine (Chap. XIII.) must of course be taken into account.

5. Another method of testing astigmatism is by means of Tweedy's optometer, of which, with the exception that the later instruments are marked in dioptries and inches instead of only in inches, he has published a description in the *Lancet*, for Oct. 28, 1876. The patient under atropine, is made artificially myopic by a convex lens, a card with fine radiating lines is gradually approached to his eye, until *one* line becomes quite distinct. The meridian at *right angles* to this line is then known to be the *least* refractive. The *concave* cylinder is found, which, with its axis at *right angles* to the line first seen, makes the line in the opposite meridian equally distinct. This shows that the latter meridian is now made as little refracting as the former. The distance at which the first line is seen indicates the kind and degree of the error for the *least* refracting meridian. A *spherical* lens correcting this is then ordered, and, in combination with it, the *concave* cylinder of the strength, and in the axis, which were found necessary to equalise the two meridians.

6. Retinoscopy. This is an exceedingly useful means of estimating astigmatism since it is an entirely objective method and places us in complete independence

of the patient. It has these advantages over the estimation of the degree of astigmatism by the ophthalmoscope that it is a much easier method to practise and that by means of it the refraction in the visual axis may be estimated. Since the refraction in the visual axis has been shewn to differ from that indicated in the optic axis. For directions see Chapter IX.

PRESBYOPIA.

Though the patient does not require glasses for distant V., he may, as age advances, find it more and more difficult to read or work without them. This defect is called *Presbyopia* and is caused chiefly by failure in the power of accommodation. A flattening of the crystalline lens sometimes accompanies it and causes some hypermetropia which has been called hypermetropia *acquisita*. *Presbyopia* is indicated by a *recession of the near point*, and it is said by Donders to have commenced when this is further from the eye than 22 cm. (9 ins.), *i.e.*, than the focal length of a lens whose refracting power is 4.50 D. ($\frac{1}{9}$). As we have already seen (Chap. IV., p. 13), such a lens would, in *Emmetropia*, bring rays from the "near point" to a focus on the retina in the absence of *all* accommodation.

The *difference*, then, between this lens and one whose focal length equals the distance of any given *receded* near point, denotes in all cases the amount of accommodation which is deficient, and the lens necessary to enable the patient again to read at 22 cm.

e.g.—With near point receded to 40 cm. (16 ins.) we

must deduct, from 4.50 D. ($\frac{1}{9}$), the lens whose focal length is 40 cm., viz., 2.50 D. ($\frac{1}{16}$), and have, as the necessary lens, + 2 D. ($\frac{1}{5}$). That is if the patient wish to read and work at 22 cm. (9 ins.)

If no accommodation remain, and either hypermetropia or hypermetropia acquisita co-exist, the strength of the correcting lens must, of course, then be *added to* 4.50 D. ($\frac{1}{9}$) in order that such a patient may read at 22 cm. (9 ins.).

The following table gives, according to Donders, the glasses necessary for Presbyopia in *Emmetropia* at different ages.

AGE.	GLASS.	
	D.	ENGLISH INCHES.
45	1	$\frac{1}{40}$
50	2	$\frac{1}{20}$
55	3	$\frac{1}{12}$
60	4	$\frac{1}{10}$
65	4.50	$\frac{1}{9}$
70	5.50	$\frac{1}{7\frac{1}{2}}$
75	6	$\frac{1}{7}$
80	7	$\frac{1}{6}$

Since these are the glasses which enable an *emmetropic* individual to read or work at 22 cm. (9 ins.) it is evident that, if either M. or H. be present, the amount of the former must be *deducted from*, and that of the latter *added to*, the lens here specified for any particular age. Though these are the glasses required in order to allow the patient

to read at 22 cm., we must test each individual case, for the various patients prefer different distances, at which to read or work.

APHAKIA

or "*absence of lens*," as after cataract operations, involves loss of accommodation. If we replace the crystalline lens by a glass in front of the eye so that parallel rays are focussed on the retina, we render the eye *practically emmetropic*. To enable such a patient to read at any specified distance, it is necessary merely to add to that glass, the lens whose focal length equals the distance in question.

e.g.—If a patient requires + 13 D. ($\frac{1}{3}$) for parallel rays and we wish him to read at a distance of 33 cm. (12 ins.) we add together 13 D. ($\frac{1}{3}$) and the lens whose focal length is 33 cm., viz. 3 D. ($\frac{1}{12}$), thus getting + 16 D. ($\frac{1}{24}$) as the necessary glass.

For cataract patients we order two pairs of spectacles. One for distant V., generally + 10 D. to + 13 D. ($\frac{1}{4}$ to $\frac{1}{3}$), and the other for reading, from + 15 D. to + 20 D. ($\frac{1}{24}$ to $\frac{1}{2}$) according to the previous state of the refraction.

We can ascertain whether the glasses obtained by patients accord with the prescription, in the following manner:—If, whilst moving a *convex* lens to and fro in front of our eye, we look at some distant object, this latter appears to move in the *opposite direction*. A contrary effect is produced with a concave lens. A concave and a convex lens of equal strength *neutralise* each other, and there is no apparent movement of the object.

If then no movement is perceptible on applying a lens of the same strength as that ordered (though of an *opposite sign*), the glasses are correct. If the lens, however, is *not* of the same strength, we can ascertain the amount of error by finding the strength of that required to effect neutralisation.

CHAPTER XIII.

ATROPINE, HOMATROPINE, HOMATROPINE AND COCAINE, ESERINE: — ALTERATIONS NECESSARY IN MEASUREMENTS MADE UNDER THEIR INFLUENCE.

ACCORDING to DEF. 3, parallel rays *should* come to a focus on the retina of an emmetropic eye when the accommodation is completely paralysed by atropine. Emmetropia, however, as thus defined, is rarely met with. An eye may have $V. = \frac{6}{8} (\frac{20}{20})$ and the distant V. may be impaired by even the *weakest* convex lens, yet, when fully under atropine, it will generally be found that a *convex* lens is *now* necessary to obtain $\frac{6}{8} (\frac{20}{20})$, *i.e.*, to bring practically parallel rays to a focus, so that a certain amount of accommodation *was* being exercised. That portion of the accommodation which can be overcome only by atropine is due to the "*tone*" of the ciliary muscle, and must always be taken into calculation when ordering glasses from a consideration of measurements of the refraction made under atropine. For *practical* purposes, we may consider that eye to be emmetropic, which, when fully under the influence of atropine, does not require a stronger convex lens than 1 D. ($\frac{1}{10}$) to perfect the focussing of parallel rays.

In *Hypermetropia* the convex lens which brings parallel rays to a focus, when the eye is fully under atropine, should *theoretically* convert it into an emmetropic eye. It is found,

however, *practically* that, owing to hypertrophy of the ciliary muscle, patients with this error of refraction find great difficulty in relaxing their accommodation to anything like its fullest extent, and consequently the lens which brings parallel rays to a focus under atropine, cannot be ordered for the same eye with its accommodation active. The increased convexity of the crystalline lens (produced by accommodation) added to the correcting lens just mentioned, renders parallel rays too convergent, and brings them to a focus in front of the retina. In Hypermetropia, therefore, it is necessary to *deduct* from the measurement made under atropine at least 1 D. ($\frac{1}{40}$), and, in the cases of children and young adults, sometimes as much as 1.50 D. ($\frac{1}{24}$) on account of the tone of the ciliary muscle.

In *Myopia* the concave lens which brings parallel rays to a focus on the retina of an eye under atropine may be found *too weak* for the same eye when *not* under atropine. The effect of the accommodation is to bring to a focus in front of the retina, the rays which, under atropine, were focussed upon it; in other words, the myopia is still sometimes uncorrected, and so requires a concave lens 0.50 D. ($\frac{1}{80}$) or 0.75 D. ($\frac{1}{50}$) stronger. But since the ciliary muscle of myopes is usually atrophied, the addition need not be equivalent to the deduction made in hypermetropia. We have, therefore, the

RULE III. That glasses ordered for permanent use must when *convex*, be 1 D. ($\frac{1}{40}$) or occasionally 1.50 D. ($\frac{1}{24}$) *weaker*, and when *concave*, 0.25 D. ($\frac{1}{160}$) to 0.75 D. ($\frac{1}{50}$) *stronger*, than is indicated by the measurements made under atropine. In M. however, this addition is seldom required.

The application of this rule may be exemplified by reference to a case of mixed astigmatism. If, under atropine, one meridian be hypermetropic, 2 D. ($\frac{1}{20}$) and the other myopic, 2 D. ($\frac{1}{20}$), the fully correcting glass would be *either* -2 D. s. $\bigcirc + 4$ D. c. ($-\frac{1}{20}$ s. $\bigcirc + \frac{1}{10}$ c.) *or* (preferably) $+ 2$ D. s. $\bigcirc - 4$ D. c. ($+\frac{1}{20}$ s. $\bigcirc - \frac{1}{10}$ c.). The glasses ordered for permanent use would be, in the first case, $- 3$ D. s. $\bigcirc + 4$ D. c. ($-\frac{1}{12}$ s. $\bigcirc + \frac{1}{10}$ c.) and in the second $+ 1$ D. s. $\bigcirc - 4$ D. c. ($+\frac{1}{40}$ s. $\bigcirc - \frac{1}{10}$ c.). In the latter example, by *deducting* 1 D. ($\frac{1}{40}$) from the convex spherical we at the same time practically add 1 D. to the strength of the concave cylinder (for it has thus *less* neutralisation to perform). While in the first case, by *adding* 1 D. to the concave spherical we lessen the strength of the convex cylinder, since it now has more to neutralise.

In *Mixed* astigmatism, then, by deducting from the strength of the spherical, we *increase* that of the cylinder: and by adding to the strength of the spherical, we *diminish* that of the cylinder.

In *Compound* astigmatism, if we add to, or deduct from, the strength of the spherical, we at the same time increase or diminish respectively the correction for each meridian.

The calculations necessary for the action of atropine may thus be made for both meridians simultaneously, by means of the *spherical* lens alone.

HOMATROPINE:

Since there is some risk in applying atropine to the eyes of people over 30 years of age, it is advisable, as shewn by Lang and Barrett in the *Medical Times*, July 18th, 1885, to apply homatropine, in such cases, or what, as they have since ascertained, is even better, a mixture of homatropine and cocaine 2 per cent. each. The accommodation and the iris are paralysed in from 30 to 60 minutes after application and the estimation of the error of Refraction can then be made. The effect, unlike that of atropine, passes away in 24 to 48 hours and consequently the risk due to the tension-raising influence of this drug is avoided.

If there is any special reason for anxiety or if the patient is particularly desirous of returning to work within that time, a few applications of a .4 per cent. eserine solution will, according to the above-mentioned observers, restore the power of accommodation in a few minutes.

The homatropine or the mixture of homatropine and cocaine should be applied in solution in castor oil.

In all estimations of errors of refraction, atropine or homatropine is absolutely necessary if perfect accuracy be sought for.

CHAPTER XIV.

AMBLYOPIA. MALINGERING AND ITS DETECTION.

It need scarcely be mentioned in conclusion that there are many cases in which, though the error of refraction has been duly diagnosed, estimated and corrected, yet there is no consequent improvement in vision. Such results may be found in old standing cases of strabismus and sometimes in eyes in which no trace of disease can be discovered. Such eyes are said to be amblyopic. Others again, without any error of refraction, may have defective vision owing to disease; with such, however, we have not here to deal. These pages have been intended merely to indicate the various modes of estimating and correcting errors of refraction; the reason why, after such correction, the vision is not improved must be ascertained by other means. Some patients, in whom we cannot find either disease or error of refraction, may simulate total or partial blindness. The latter may often be detected by holding in front of their eyes different pairs of convex, neutralised by their corresponding concave lenses. With these, thinking they are being aided, such patients may frequently be persuaded to read perfectly. Another method of detecting simulation, especially of *one* eye, is to hold in front of the *blind* (?) eye, a prism with its base out or in, when if there be an attempt (detected by the involuntary movement of the eye) to fuse the *double* images, it

proves sight present in the eye in question. Another method is to place before one eye a red, and before the other a green, glass, and then to ask the patient to read some green and red letters. If he can read both the green and the red letters he must of course see with both eyes.

If some such order of procedure, as indicated in these pages, were adopted, one would not, as is now frequently the case, see a beginner fall into the error of supposing a patient myopic because he can read $\frac{6}{8}$ ($\frac{20}{20}$) as well with, as without, a *concave* lens ; or that hypermetropia is present because the patient can read Sn. 1 with *convex* glasses.

Some of the statements may have appeared somewhat too general but it has been thought advisable not to confuse the beginner by enumerating all the possible and minor exceptions to the general rules.



D=0,5

J. 1.

I now stood a calm spectator of the flames, and after some time began to perceive that my arm to the shoulder was scorched in a terrible manner. It was, therefore, out of my power to give my son any assistance, either in attempting to save our goods, or preventing the flames spreading to our corn.

D = 0,6

J. 2.

Sometimes to give a variety to our amusements, the girls sang to the guitar; and while they thus formed a little concert, my wife and I would stroll down the sloping field, that was fragrant with wild flowers, and enjoy the breeze that wafted both health and harmony.

D=0,8

J. 4.

At last a young gentleman, of a more genteel appearance than the rest, came forward, and for a while regarding us, stopped short, and giving his horse to a servant who attended, approached us with a careless superior air.

D=1

J. 6.

The family seemed earnest to please him; my girls entertained him with topics they thought most modern, while Moses gave him a question or two from the ancients, for which he had the satisfaction of being laughed at.

D=1,25

J. 8.

Such was the power of fortune and fine clothes, that he found no second repulse. In course of conversation

D=1,5

J. 10.

He let us know that he was the owner of the splendid picturesque estate that lay round us.

D=1,75

J. 12.

At these times our little ones always read short tales for our amusement.

D=2,25

J. 14.

**For in this way you may
always damp our ardour.**

D=3

J. 15.

I saw no one there.

D=4

J. 16.

For the ensuing

PUBLISHED BY
CURRY & PAXTON, OPHTHALMIC OPTICIANS,
195, GREAT PORTLAND STREET, LONDON, W.
LIVERPOOL AND BRISTOL.

D=60 (200)

IR

D=36 (120)

ET

D=24 (80)

I A V

D=18 (60)

R T B

D=12 (40)

D F O E

D=9 (30)

C L Z T O


D=6 (20)

L T R F P

D=5 (16)

A P O R F D

ASTIGMATIC FAN.



CATALOGUE OF WORKS

PUBLISHED BY

H. K. LEWIS

136 GOWER STREET, LONDON, W.C.

Established 1844.

SIR WILLIAM AITKEN, KNT., M.D., F.R.S.

Professor of Pathology in the Army Medical School.

ON THE ANIMAL ALKALOIDS, THE PTOMAINES, LEUCOMAINES, AND EXTRACTIVES IN THEIR PATHOLOGICAL RELATIONS. Second edition, Crown 8vo, 3s. 6d.

E. CRESSWELL BABER, M.B. LOND.

Surgeon to the Brighton and Sussex Throat and Ear Dispensary.

A GUIDE TO THE EXAMINATION OF THE NOSE, WITH REMARKS ON THE DIAGNOSIS OF DISEASES OF THE NASAL CAVITIES. With Illustrations, small 8vo, 5s. 6d.

JAMES B. BALL, M.D. LOND., M.R.C.P.

Physician to the Department for Diseases of the Throat and Nose, and Senior Assistant Physician, West London Hospital.

I.

A HANDBOOK OF DISEASES OF THE NOSE AND NASO-PHARYNX. Large post 8vo, with illustrations, 6s.

II.

INTUBATION OF THE LARYNX. With Illustrations, Demy 8vo, 2s. 6d.

G. GRANVILLE BANTOCK, M.D., F.R.C.S. EDIN.

Surgeon to the Samaritan Free Hospital for Women and Children.

I.

ON THE USE AND ABUSE OF PESSARIES. Second Edit. with Illustrations, 8vo, 5s.

II.

ON THE TREATMENT OF RUPTURE OF THE FEMALE PERINEUM IMMEDIATE AND REMOTE. Second Edition, with Illustrations, 8vo, 3s. 6d.

III.

A PLEA FOR EARLY OVARIOTOMY. Demy 8vo, 2s.

ARTHUR E. J. BARKER, F.R.C.S.

Hunterian Professor of Surgery and Pathology; Surgeon to University College Hospital.

HUNTERIAN LECTURES ON INTRA-CRANIAL INFLAMMATIONS STARTING IN THE TEMPORAL BONE THEIR COMPLICATIONS AND TREATMENT. 8vo, 3s. nett.

DR. FERBER.

MODEL DIAGRAM OF THE ORGANS IN THE THORAX AND UPPER PART OF THE ABDOMEN. With Letter-press Description. In 4to, coloured, 5s.

J. MAGEE FINNY, M.D. DUBL.

King's Professor of Practice of Medicine in School of Physic, Ireland, &c.

NOTES ON THE PHYSICAL DIAGNOSIS OF LUNG DISEASES. 32mo, 1s. 6d. [Now ready.]

AUSTIN FLINT, M.D., LL.D.

Professor of Physiology and Physiological Anatomy in the Bellevue Hospital Medical College, New York; visiting Physician to the Bellevue Hospital, &c.

I.

A TEXT-BOOK OF HUMAN PHYSIOLOGY. Fourth edition, Illustrated by plates, and 316 wood engravings, large 8vo, 25s.

II.

THE PHYSIOLOGY OF THE SPECIAL SENSES AND GENERATION; (Being Vol. V. of the Physiology of Man). Roy. 8vo, 18s.

J. MILNER FOTHERGILL, M.D., M.R.C.P.

Late Physician to the City of London Hospital for Diseases of the Chest, Victoria Park, &c.

I.

A MANUAL OF DIETETICS. Large 8vo, 10s. 6d.

II.

THE HEART AND ITS DISEASES, WITH THEIR TREATMENT; INCLUDING THE GOUTY HEART. Second Edition, entirely re-written, copiously illustrated with woodcuts and lithographic plates. 8vo, 16s.

III.

INDIGESTION AND BILIOUSNESS. Second Edition, post 8vo, 7s. 6d.

IV.

GOUT IN ITS PROTEAN ASPECTS. Post 8vo, 7s. 6d.

V.

THE TOWN DWELLER: His Needs and His Wants. With an Introduction by B. W. RICHARDSON, M.D., LL.D., F.R.S. Post 8vo, 3s. 6d. [Now ready.]

FORTESCUE FOX, M.D. LOND.

Fellow of the Medical Society of London.

STRATHPEFFER SPA: Its Climate and Waters. With OBSERVATIONS HISTORICAL, MEDICAL, AND GENERAL, DESCRIPTIVE OF THE VICINITY. Crown 8vo, with Map and Illustrations, 2s. 6d., nett.

ERNEST FRANCIS, F.C.S.

Demonstrator of Practical Chemistry, Charing Cross Hospital.

PRACTICAL EXAMPLES IN QUANTITATIVE ANALYSIS, forming a Concise Guide to the Analysis of Water, &c. Illustrated, fcap. 8vo, 2s. 6d.

JOHN HENRY GARRETT, M.D.

Licentiate in Sanitary Science and Diplomate in Public Health, Universities of Durham and Cambridge, &c.

THE ACTION OF WATER ON LEAD; being an inquiry into the Cause and Mode of the Action and its Prevention. Crown 8vo, 4s. 6d.

ALFRED W. GERRARD, F.C.S.

Examiner to the Pharmaceutical Society; Teacher of Materia Medica and Pharmacy at University College Hospital.

ELEMENTS OF MATERIA MEDICA AND PHARMACY. Crown 8vo, 8s. 6d.

NEW OFFICIAL REMEDIES, B.P., 1890. Supplement to the above. Crown 8vo, 1s.

HENEAGE GIBBES, M.D.

Lecturer on Physiology and on Normal and Morbid Histology in the Medical School of Westminster Hospital; etc.

PRACTICAL HISTOLOGY AND PATHOLOGY. Third Edition, revised and enlarged, crown 8vo, 6s.

C. A. GORDON, M.D., C.B.

Deputy Inspector General of Hospitals, Army Medical Department.

REMARKS ON ARMY SURGEONS AND THEIR WORKS. Demy 8vo, 5s.

JOHN GORHAM, M.R.C.S.

TOOTH EXTRACTION: a Manual on the proper mode of extracting Teeth. Third Edition, fcap. 8vo, 1s. 6d. [Now ready

GEORGE M. GOULD, B.A., M.D.

Ophthalmic Surgeon to the Philadelphia Hospital, etc.

A NEW MEDICAL DICTIONARY: including all the words and phrases used in Medicine, with their proper pronunciation and definitions. 8vo, 16s.

W. R. GOWERS, M.D., F.R.C.P., M.R.C.S.

Physician to University College Hospital, &c.

DIAGRAMS FOR THE RECORD OF PHYSICAL SIGNS. In books of 12 sets of figures, 1s. Ditto, unbound, 1s.

J. B. GRESSWELL, M.R.C.V.S.

Provincial Veterinary Surgeon to the Royal Agricultural Society.

VETERINARY PHARMACOLOGY AND THERAPEUTICS. With an Index of Diseases and Remedies. Fcap. 8vo, 5s.

SAMUEL D. GROSS, M.D., LL.D., D.C.L. OXON.

Professor of Surgery in the Jefferson Medical College of Philadelphia.

A PRACTICAL TREATISE ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE URINARY BLADDER, THE PROSTATE GLAND, AND THE URETHRA. Third Edition, revised and edited by S. W. GROSS, A.M., M.D., Surgeon to the Philadelphia Hospital. Illustrated by 170 engravings, 8vo, 18s.

SAMUEL W. GROSS, A.M., M.D.

Surgeon to, and Lecturer on Clinical Surgery in, the Jefferson Medical College Hospital, and the Philadelphia Hospital, &c.

- A PRACTICAL TREATISE ON TUMOURS OF THE MAMMARY GLAND:** embracing their Histology, Pathology, Diagnosis, and Treatment. With Illustrations, 8vo, 10s. 6d.

DR. JOSEF GRUBER.

Professor of Otology in the Imperial Royal University of Vienna, etc.

- A TEXT-BOOK OF THE DISEASES OF THE EAR.**
Translated from the second German edition by special permission of the Author, and Edited by EDWARD LAW, M.D., C.M. EDIN., M.R.C.S. ENG., Surgeon to the London Throat Hospital for Diseases of the Throat, Nose and Ear; and by COLEMAN JEWELL, M.B. LOND., M.R.C.S. ENG., late Physician and Pathologist to the London Throat Hospital. With 150 Illustrations, and 70 coloured figures on 2 lithographic plates, Royal 8vo, 24s. [Just Published.]

ALLAN McLANE HAMILTON, M.D.

- THE MODERN TREATMENT OF HEADACHES.**
Square 16mo, 2s. 6d.

WILLIAM A. HAMMOND, M.D.

Professor of Mental and Nervous Diseases in the Medical Department of the University of the City of New York, &c.

- SPIRITUALISM AND ALLIED CAUSES AND CONDITIONS OF NERVOUS DERANGEMENT.** With Illustrations, post 8vo, 8s. 6d.

ALEXANDER HARVEY, M.D.

Late Emeritus Professor of Materia Medica in the University of Aberdeen, &c.,

AND

ALEXANDER DYCE DAVIDSON, M.D., F.R.S. EDIN.

Late Regius Professor of Materia Medica in the University of Aberdeen.

- SYLLABUS OF MATERIA MEDICA FOR THE USE OF STUDENTS, TEACHERS AND PRACTITIONERS.** Based on the relative values of articles and preparations in the British Pharmacopœia. Ninth edition, 32mo, 1s. 6d.

K. M. HEANLEY.

Matron of Boston Cottage Hospital.

- A MANUAL OF URINE TESTING.** Compiled for the use of Matrons, Nurses, and Probationers. Post 8vo, 1s. 6d.

C. HIGGENS, F.R.C.S.

Ophthalmic Surgeon to Guy's Hospital; Lecturer on Ophthalmology at Guy's Hospital Medical School.

- MANUAL OF OPHTHALMIC PRACTICE.**

Crown 8vo, illustrations, 6s.

[LEWIS'S PRACTICAL SERIES.]

BERKELEY HILL, M.B. LOND., F.R.C.S.

Professor of Clinical Surgery in University College; Surgeon to University College Hospital and to the Lock Hospital.

THE ESSENTIALS OF BANDAGING. With directions for Managing Fractures and Dislocations; for administering Ether and Chloroform; and for using other Surgical Apparatus; with a Chapter on Surgical Landmarks. Sixth Edition, revised and enlarged, Illustrated by 144 Wood Engravings, crown 8vo, 5s.

BERKELEY HILL, M.B. LOND., F.R.C.S.

Professor of Clinical Surgery in University College; Surgeon to University College Hospital and to the Lock Hospital.

AND

ARTHUR COOPER, L.R.C.P., M.R.C.S.

Surgeon to the Westminster General Dispensary.

I

SYPHILIS AND LOCAL CONTAGIOUS DISORDERS.
Second edition, entirely re-written, royal 8vo, 18s.

II.

THE STUDENT'S MANUAL OF VENEREAL DISEASES. Being a Concise Description of those Affections and of their Treatment. Fourth edition, post 8vo, 2s. 6d.

PROCTER S. HUTCHINSON, M.R.C.S.

Assistant Surgeon to the Hospital for Diseases of the Throat.

A MANUAL OF DISEASES OF THE NOSE AND THROAT; including the Nose, Naso-pharynx, Pharynx, and Larynx. With Illustrations, crown 8vo, 3s. 6d. *[Now ready.]*

C. R. ILLINGWORTH, M.D. ED., M.R.C.S.

THE ABORTIVE TREATMENT OF SPECIFIC FEBRILE DISORDERS BY THE BINIODIDE OF MERCURY.
Crown 8vo, 3s. 6d.

SIR W. JENNER, Bart., M.D.

Physician in Ordinary to H.M. the Queen, and to H.R.H. the Prince of Wales.

THE PRACTICAL MEDICINE OF TO-DAY: Two Addresses delivered before the British Medical Association, and the Epidemiological Society, (1869). Small 8vo, 1s. 6d.

GEORGE LINDSAY JOHNSON, M.A., M.B., B.C. CANTAB.
Clinical Assistant, late House Surgeon and Chloroformist, Royal Westminster Ophthalmic Hospital, &c.

A NEW METHOD OF TREATING CHRONIC GLAUCOMA, based on Recent Researches into its Pathology. With Illustrations and coloured frontispiece, demy 8vo, 3s. 6d.

JOHN M KEATING,
Fellow of the College of Physicians, Philadelphia, &c.
 AND
 HENRY HAMILTON.
POCKET MEDICAL LEXICON.
 32mo, 3s. nett.

NORMAN KERR, M.D., F.L.S.
President of the Society for the Study of Inebriety; Consulting Physician, Dalrymple Home for Inebriates, &c.
INEBRIETY: its Etiology, Pathology, Treatment, and Jurisprudence. Second edition, Crown 8vo, 12s. 6d.

NORMAN W. KINGSLEY, M.D.S., D.D.S.
President of the Board of Censors of the State of New York; Member of the American Academy of Dental Science, &c.
A TREATISE ON ORAL DEFORMITIES AS A BRANCH OF MECHANICAL SURGERY. With over 350 Illustrations, 8vo, 16s.

F. CHARLES LARKIN, F.R.C.S. ENG.
Assistant Lecturer in Physiology in University College, Liverpool.
 AND
 RANDLE LEIGH, M.B., B.SC. LOND.
Senior Demonstrator of Physiology in University College, Liverpool.
OUTLINES OF PRACTICAL PHYSIOLOGICAL CHEMISTRY. With Illustrations, crown 8vo. [Just ready.]

J. WICKHAM LEGG, F.R.C.P.
Assistant Physician to Saint Bartholomew's Hospital, and Lecturer on Pathological Anatomy in the Medical School.
 I.
ON THE BILE, JAUNDICE, AND BILIOUS DISEASES.
 With Illustrations in chromo-lithography, 719 pages, roy. 8vo, 25s.
 II.
A GUIDE TO THE EXAMINATION OF THE URINE;
 intended chiefly for Clinical Clerks and Students. Sixth Edition, revised and enlarged, with Illustrations, fcap. 8vo, 2s. 6d.
 III.
A TREATISE ON HÆMOPHILIA, SOMETIMES CALLED THE HEREDITARY HÆMORRHAGIC DIATHESIS.
 Fcap. 4to, 7s. 6d.

ARTHUR H. N. LEWERS, M.D. LOND., M.R.C.P. LOND.
Assistant Obstetric Physician to the London Hospital; Examiner in Midwifery and Diseases of Women to the Society of Apothecaries of London, &c.
A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN. Third edition, Illustrations, crown 8vo, 10s. 6d. [Now ready.]
 [LEWIS'S PRACTICAL SERIES.]

LEWIS'S POCKET CASE BOOK FOR PRACTITIONERS AND STUDENTS. Designed by A. T. BRAND, M.D. Roan, with pencil, 3s. 6d. *nett.*

LEWIS'S POCKET MEDICAL VOCABULARY.
Second Edition, thoroughly revised, 32mo, roan, 3s. 6d. [*Just ready.*]

T. R. LEWIS, M.B., F.R.S. ELECT, ETC.
Late Fellow of the Calcutta University, Surgeon-Major Army Medical Staff, &c.
PHYSIOLOGICAL AND PATHOLOGICAL RESEARCHES. Arranged and edited by SIR WM. AITKEN, M.D., F.R.S., G. E. DOBSON, M.B., F.R.S., and A. E. BROWN, B.Sc. Crown 4to, portrait, 5 maps, 43 plates including 15 chromo-lithographs, and 67 wood engravings, 30s. *nett.*

* * A few copies only of this work remain for sale.

C. B. LOCKWOOD, F.R.C.S.
Hunterian Professor, Royal College of Surgeons of England; Surgeon to the Great Northern Hospital; Senior Demonstrator of Anatomy and Operative Surgery in St. Bartholomew's Hospital.

HUNTERIAN LECTURES ON THE MORBID ANATOMY, PATHOLOGY AND TREATMENT OF HERNIA. Demy 8vo, 36 illustrations, 5s.

J. S. LOMBARD, M.D.
Formerly Assistant Professor of Physiology in Harvard College.
I.
EXPERIMENTAL RESEARCHES ON THE REGIONAL TEMPERATURE OF THE HEAD, under Conditions of Rest, Intellectual Activity, and Emotion. With Illustrations, 8vo, 8s.
II.
ON THE NORMAL TEMPERATURE OF THE HEAD.
8vo, 5s.

WILLIAM THOMPSON LUSK, M.A., M.D.
Professor of Obstetrics and Diseases of Women in the Bellevue Hospital Medical College, &c
THE SCIENCE AND ART OF MIDWIFERY.
Third Edition, with numerous Illustrations, 8vo, 18s.

A. W. MACFARLANE, M.D., F.R.C.P. EDIN.
Examiner in Medical Jurisprudence in the University of Glasgow; Honorary Consulting Physician (late Physician) Kilmarnock Infirmary
INSOMNIA AND ITS THERAPEUTICS.
Medium 8vo, 12s. 6d.

SURGEON-MAJOR C. J. McNALLY, M.D., D.P.H. CAMB.
Fellow of the Madras University; Professor of Chemistry, Madras Medical College.
THE ELEMENTS OF SANITARY SCIENCE.
Plates, Demy 8vo, 8s. 6d.

RAWDON MACNAMARA.

Professor of Materia Medica, Royal College of Surgeons, Ireland; Senior Surgeon to the Westmoreland (Lock) Government Hospital; Surgeon to the Meath Hospital, &c.

**AN INTRODUCTION TO THE STUDY OF THE
BRITISH PHARMACOPŒIA.** Demy 32mo, 1s. 6d. [*Just published.*]

JOHN MACPHERSON, M.D.

*Inspector-General of Hospitals H.M. Bengal Army (Retired).
Author of "Cholera in its Home," &c.*

I.

**ANNALS OF CHOLERA FROM THE EARLIEST
PERIODS TO THE YEAR 1817.** With a map. Demy 8vo, 7s. 6d.

II.

**BATH, CONTREXEVILLE, AND THE LIME SUL-
PHATED WATERS.** Crown 8vo, 2s. 6d.

A. COWLEY MALLEY, B.A., M.B., B.CH. T.C.D.

PHOTO-MICROGRAPHY; including a description of the Wet Collodion and Gelatino-Bromide Processes, together with the best methods of Mounting and Preparing Microscopic Objects for Photo-Micrography. Second Edition, with Photographs and Illustrations, crown 8vo, 7s. 6d.

PATRICK MANSON, M.D., C.M.

**THE FILARIA SANGUINIS HOMINIS; AND CER-
TAIN NEW FORMS OF PARASITIC DISEASE IN INDIA,
CHINA, AND WARM COUNTRIES.** Illustrated with Plates and Charts. 8vo, 10s. 6d.

JEFFERY A. MARSTON, M.D., C.B., F.R.C.S., M.R.C.P. LOND.
Surgeon General Medical Staff (Retired).

**NOTES ON TYPHOID FEVER: Tropical Life and its
Sequelæ.** Crown 8vo, 3s. 6d. [*Now ready.*]

PROFESSOR MARTIN.

**MARTIN'S ATLAS OF OBSTETRICS AND GYNÆCO-
LOGY.** Edited by A. MARTIN, Docent in the University of Berlin, Translated and edited with additions by FANCOURT BARNES, M.D., M.R.C.P., Physician to the Chelsea Hospital for Women; Obstetric Physician to the Great Northern Hospital; and to the Royal Maternity Charity of London, &c. Medium 4to, Morocco half bound, 31s. 6d. *nett.*

EDWARD MARTIN, A.M., M.D.
**MINOR SURGERY AND BANDAGING WITH AN
APPENDIX ON VENEREAL DISEASES.** Crown 8vo, 82 Illus-
trations, 4s.

WILLIAM MARTINDALE, F.C.S.
*Late Examiner of the Pharmaceutical Society, and late Teacher of Pharmacy and Demon-
strator of Materia Medica at University College,*

AND

W. WYNN WESTCOTT, M.B. LOND.
Deputy Coroner for Central Middlesex.

THE EXTRA PHARMACOPŒIA with the additions in-
troduced into the British Pharmacopœia, 1885 and 1890, with Medical
References, and a Therapeutic Index of Diseases and Symptoms. Sixth
Edition, limp roan, med. 24mo, 7s. 6d. [Now ready.]

WILLIAM MARTINDALE, F.C.S.
Late Examiner of the Pharmaceutical Society, &c.
COCA, COCAINE, AND ITS SALTS: their History,
Medical and Economic Uses, and Medicinal Preparations. Fcap. 8vo, 2s.

MATERIA MEDICA LABELS.

Adapted for Public and Private Collections. Compiled from the British
Pharmacopœia of 1885, with the additions of 1890. The Labels are ar-
ranged in Two Divisions:—

Division I.—Comprises, with few exceptions, Substances of Organ-
ized Structure, obtained from the Vegetable and Animal King-
doms.

Division II.—Comprises Chemical Materia Medica, including Alco-
hols, Alkaloids, Sugars, and Neutral Bodies.

On plain paper, 10s. 6d. *nett.* On gummed paper, 12s. 6d. *nett.*

The 24 additional Labels of 1890 only, 1s. *nett.*

. Specimens of the Labels, of which there are over 470, will be sent on application.

S. E. MAUNSELL, L.R.C.S.I.
Surgeon-Major, Medical Staff.
NOTES OF MEDICAL EXPERIENCES IN INDIA
PRINCIPALLY WITH REFERENCE TO DISEASES OF THE
EYE. With Map, post 8vo, 3s. 6d.

J. F. MEIGS, M.D.
Consulting Physician to the Children's Hospital, Philadelphia,
AND
W. PEPPER, M.D.
Lecturer on Clinical Medicine in the University of Pennsylvania.
**A PRACTICAL TREATISE ON THE DISEASES OF
CHILDREN.** Seventh Edition, revised and enlarged, roy. 8vo, 28s.

WM. JULIUS MICKLE, M.D., F.R.C.P. LOND.
Medical Superintendent, Grove Hall Asylum, London, &c.

I.
GENERAL PARALYSIS OF THE INSANE.

Second Edition, enlarged and rewritten, 8vo, 14s.

II.
**ON INSANITY IN RELATION TO CARDIAC AND
 AORTIC DISEASE AND PHTHISIS.** Crown 8vo, 3s. 6d.

KENNETH W. MILLICAN, B.A. CANTAB., M.R.C.S.
THE EVOLUTION OF MORBID GERMS: A Contribution to Transcendental Pathology. Cr. 8vo, 3s. 6d.

ANGEL MONEY, M.D. LOND., F.R.C.P.
*Assistant Physician to University College Hospital, and to the Hospital for Sick Children
 Great Ormond Street; Assistant Professor of Clinical Medicine in University
 College, London, &c.*

I.
**THE STUDENT'S TEXTBOOK OF THE PRACTICE
 OF MEDICINE.** Fcap. 8vo, 6s. 6d.

II.
**TREATMENT OF DISEASE IN CHILDREN: EM-
 BODYING THE OUTLINES OF DIAGNOSIS AND THE
 CHIEF PATHOLOGICAL DIFFERENCES BETWEEN CHILD-
 REN AND ADULTS.** Second edition, crown 8vo, 10s. 6d.
 [LEWIS'S PRACTICAL SERIES.]

E. A. MORSHEAD, M.R.C.S., L.R.C.P.
Assistant to the Professor of Medicine in University College, London.
**TABLES OF THE PHYSIOLOGICAL ACTION OF
 DRUGS.** Fcap. 8vo, 1s.

A. STANFORD MORTON, M.B., F.R.C.S. ENG.
Surgeon to the Royal South London Ophthalmic Hospital.
**REFRACTION OF THE EYE: Its Diagnosis, and the
 Correction of its Errors.** Fourth Edition, with Illustrations, small 8vo,
 3s. 6d. [Just ready.]

C. W. MANSELL MOULLIN, M.A., M.D. OXON., F.R.C.S. ENG.
*Assistant Surgeon and Senior Demonstrator of Anatomy at the London Hospital; formerly
 Radcliffe Travelling Fellow and Fellow of Pembroke College, Oxford.*
**SPRAINS; THEIR CONSEQUENCES AND TREAT-
 MENT.** Crown 8vo, 5s.

PAUL F. MUNDE, M.D.
*Professor of Gynecology at the New York Polyclinic; President of the New York Obstetrical
 Society and Vice-President of the British Gynecological Society, &c.*
**THE MANAGEMENT OF PREGNANCY, PARTURI-
 TION, AND THE PUERPERAL STATE.** Second edition, square
 8vo, 3s. 6d.

WILLIAM MURRAY, M.D., F.R.C.P. LOND.

Consulting Physician to the Children's Hospital, Newcastle-on-Tyne, &c.

**ILLUSTRATIONS OF THE INDUCTIVE METHOD IN
MEDICINE.** Crown 8vo, 3s. 6d. [Just ready.]

WILLIAM MURRELL, M.D., F.R.C.P.

*Lecturer on Pharmacology and Therapeutics at Westminster Hospital; late Examiner in
Materia Medica to the Royal College of Physicians of London, etc.*

I.

**MASSOTHERAPEUTICS, OR MASSAGE AS A MODE
OF TREATMENT.** Fifth edition, with Illustrations, crown 8vo,
4s. 6d. [Now ready.]

II.

WHAT TO DO IN CASES OF POISONING.
Sixth edition, royal 32mo, 3s. 6d.

III.

**NITRO-GLYCERINE AS A REMEDY FOR ANGINA
PECTORIS.** Crown 8vo, 3s. 6d.

IV.

CHRONIC BRONCHITIS AND ITS TREATMENT.
Crown 8vo, 3s. 6d.

DR. FELIX von NIEMEYER.

*Late Professor of Pathology and Therapeutics; Director of the Medical Clinic of the
University of Tübingen.*

**A TEXT-BOOK OF PRACTICAL MEDICINE, WITH
PARTICULAR REFERENCE TO PHYSIOLOGY AND PATHO-
LOGICAL ANATOMY.** Translated from the Eighth German Edi-
tion by special permission of the Author, by GEORGE H. HUMPHERY,
M.D., and CHARLES E. HACKLEY, M.D. Revised edition, 2 vols.
large 8vo, 36s.

GEORGE OLIVER, M.D., F.R.C.P.

I.

THE HARROGATE WATERS: Data Chemical and Therapeu-
tical, with notes on the Climate of Harrogate. Addressed to the
Medical Profession. Crown 8vo, with Map of the Wells, 3s. 6d.

II.

ON BEDSIDE URINE TESTING: a Clinical Guide to the
Observation of Urine in the course of Work. Fourth Edition, fcap.
8vo, 3s. 6d.

SAMUEL OSBORN, F.R.C.S.

Surgeon to the Hospital for Women, Soho Square; Surgeon Royal Naval Artillery Volunteers.

I.
AMBULANCE LECTURES: FIRST AID. Second edition,
with Illustrations, fcap. 8vo, 1s. 6d.

II.
**AMBULANCE LECTURES: HOME NURSING AND
HYGIENE.** Second edition, With Illustrations, fcap. 8vo, 2s.
[Just published.]

WILLIAM OSLER, M.D., F.R.C.P. LOND.

Professor of Clinical Medicine in the University of Pennsylvania, &c.

THE CEREBRAL PALSIES OF CHILDREN. A Clinical
Study from the Infirmary for Nervous Diseases, Philadelphia. Demy
8vo, 5s. [Just Published.]

K. W. OSTROM.

*Instructor in Massage and Swedish Movements in the Hospital of the University of
Pennsylvania.*

**MASSAGE AND THE ORIGINAL SWEDISH MOVE-
MENTS.** With illustrations, 12mo, 2s. 6d. nett. [Now ready:]

ROBERT W. PARKER.

*Senior Surgeon to the East London Hospital for Children; Surgeon to the German
Hospital.*

I.
**DIPHTHERIA: ITS NATURE AND TREATMENT,
WITH SPECIAL REFERENCE TO THE OPERATION, AFTER-
TREATMENT AND COMPLICATIONS OF TRACHEOTOMY.**
Third Edition, with Illustrations, 8vo, 6s. [Now ready.]

II.
**CONGENITAL CLUB-FOOT; ITS NATURE AND
TREATMENT.** With special reference to the subcutaneous division
of Tarsal Ligaments. 8vo, 7s. 6d.

LOUIS C. PARKES, M.D., D.P.H. LOND. UNIV.

*Fellow of the Sanitary Institute, and Member of the Board of Examiners; Assistant Pro-
fessor of Hygiene and Public Health at University College, London, &c.*

HYGIENE AND PUBLIC HEALTH. Second edition, with
numerous Illustrations, crown 8vo, 9s. [Just Published.]
[LEWIS'S PRACTICAL SERIES.]

JOHN S. PARRY, M.D.

*Obstetrician to the Philadelphia Hospital, Vice-President of the Obstetrical and Pathologi-
cal Societies of Philadelphia, &c.*

**EXTRA-UTERINE PREGNANCY; Its Causes, Species,
Pathological Anatomy, Clinical History, Diagnosis, Prognosis and
Treatment.** 8vo, 8s.

THEOPHILUS PARVIN, M.D.

Professor of Obstetrics and Diseases of Women and Children at the Jefferson Medical School.

LECTURES ON OBSTETRIC NURSING, Delivered at the Training School for Nurses of the Philadelphia Hospital. Post 8vo, 2s. 6d.

E. RANDOLPH PEASLEE, M.D., LL.D.

Late Professor of Gynæcology in the Medical Department of Dartmouth College; President of New York Academy of Medicine, &c., &c.

OVARIAN TUMOURS: Their Pathology, Diagnosis, and Treatment, especially by Ovariectomy. Illustrations, roy. 8vo, 16s.

HENRY G. PIFFARD, A.M., M.D.

Clinical Professor of Dermatology, University of the City of New York; Surgeon in Charge of the New York Dispensary for Diseases of the Skin, &c.

A PRACTICAL TREATISE ON DISEASES OF THE SKIN. With 50 full page Original Plates and 33 Illustrations in the Text, 4to, 52s. 6d. *nett.* [Just published.]

G. V. POORE, M.D., F.R.C.P.

Professor of Medical Jurisprudence, University College; Assistant Physician to, and Physician in charge of the Throat Department of, University College Hospital.

LECTURES ON THE PHYSICAL EXAMINATION OF THE MOUTH AND THROAT. With an Appendix of Cases. 8vo, 3s. 6d.

R. DOUGLAS POWELL, M.D., F.R.C.P., M.R.C.S.

Physician in Ordinary to H.M. the Queen; Physician to the Middlesex Hospital and Physician to the Hospital for Consumption and Diseases of the Chest at Brompton.

I.
DISEASES OF THE LUNGS AND PLEURÆ, INCLUDING CONSUMPTION. Third edition, entirely rewritten and enlarged. With coloured plates and wood engravings, 8vo, 16s.

II.
TABLE OF PHYSICAL EXAMINATION OF THE LUNGS—with Note on International Nomenclature of Physical Signs (reprinted from above). On one sheet, 6d.

URBAN PRITCHARD, M.D. EDIN., F.R.C.S. ENG.

Professor of Aural Surgery at King's College, London; Aural Surgeon to King's College Hospital; Senior Surgeon to the Royal Ear Hospital.

HANDBOOK OF DISEASES OF THE EAR FOR THE USE OF STUDENTS AND PRACTITIONERS. Second edition, With Illustrations, crown 8vo, 5s. [LEWIS'S PRACTICAL SERIES.]

CHARLES W. PURDY, M.D. (QUEEN'S UNIV.)

Professor of Genito-Urinary and Renal Diseases in the Chicago Polyclinic, &c., &c.

BRIGHT'S DISEASE AND THE ALLIED AFFECTIONS OF THE KIDNEYS. With Illustrations, large 8vo, 8s. 6d.

DR. THEODOR PUSCHMANN.

Public Professor in Ordinary at the University of Vienna.

HISTORY OF MEDICAL EDUCATION FROM THE MOST REMOTE TO THE MOST RECENT TIMES. Translated by EVAN H. HARE, M.A. OXON., F.R.C.S. ENG., L.S.A. Demy 8vo. [In the press.]

CHARLES HENRY RALFE, M.A., M.D. CANTAB., F.R.C.P. LOND.
Assistant Physician to the London Hospital; Examiner in Medicine to the University of Durham, &c., &c.

A PRACTICAL TREATISE ON DISEASES OF THE KIDNEYS AND URINARY DERANGEMENTS. With Illustrations, crown 8vo, 10s. 6d. [LEWIS'S PRACTICAL SERIES.]

FRANCIS H. RANKIN, M.D.

President of the New York Medical Society.

HYGIENE OF CHILDHOOD. Suggestions for the care of Children after the Period of Infancy to the completion of Puberty. Crown 8vo, 3s.

AMBROSE L. RANNEY, A.M., M.D.

Professor of the Anatomy and Physiology of the Nervous System in the New York Post-Graduate Medical School and Hospital, &c.

THE APPLIED ANATOMY OF THE NERVOUS SYSTEM. Second edition, 238 Illustrations, large 8vo, 21s.

H. A. REEVES, F.R.C.S. EDIN.

Senior Assistant Surgeon and Teacher of Practical Surgery at the London Hospital; Surgeon to the Royal Orthopædic Hospital.

BODILY DEFORMITIES AND THEIR TREATMENT: A HANDBOOK OF PRACTICAL ORTHOPÆDICS. Illustrations, crown 8vo, 8s. 6d. [LEWIS'S PRACTICAL SERIES.]

RALPH RICHARDSON, M.A., M.D.

Fellow of the College of Physicians, Edinburgh.

ON THE NATURE OF LIFE: An Introductory Chapter to Pathology. Second edition, revised and enlarged. Fcap. 4to, 10s. 6d.

W. RICHARDSON, M.A., M.D., M.R.C.P.

REMARKS ON DIABETES, ESPECIALLY IN REFERENCE TO TREATMENT. Demy 8vo, 4s. 6d.

SAMUEL RIDEAL, D.SC. (LOND.), F.I.C., F.C.S., F.G.S.

Fellow of University College, London.

PRACTICAL ORGANIC CHEMISTRY; The Detection and Properties of some of the more important Organic Compounds. 12mo, 2s. 6d.

II.

PRACTICAL CHEMISTRY FOR MEDICAL STUDENTS, required at the First Examination of the Conjoint Examining Board in England. Foolscap 8vo 2s. [Just published.]

E. A. RIDSDALE.

Associate of the Royal School of Mines.

COSMIC EVOLUTION ; being Speculations on the Origin
of our Environment. Fcap. 8vo, 3s.

SYDNEY RINGER, M.D., F.R.S.

*Professor of the Principles and Practice of Medicine in University College ; Physician to,
and Professor of Clinical Medicine in, University College Hospital.*

I.

A HANDBOOK OF THERAPEUTICS. Twelfth Edition
thoroughly revised, 8vo, 15s.

II.

**ON THE TEMPERATURE OF THE BODY AS
A MEANS OF DIAGNOSIS AND PROGNOSIS IN PHTHISIS.**
Second edition, small 8vo, 2s. 6d.

FREDERICK T. ROBERTS, M.D., B.SC., F.R.C.P.

*Examiner in Medicine at the University of London and for the Conjoint Board ; Professor
of Materia Medica and Therapeutics and of Clinical Medicine in University
College ; Physician to University College Hospital ; Physician to
Brompton Consumption Hospital, &c.*

I.

**A HANDBOOK OF THE THEORY AND PRACTICE
OF MEDICINE.** Eighth edition, with Illustrations, in one volume,
large 8vo, 21s. [Just published.]

II.

THE OFFICINAL MATERIA MEDICA.

Second edition, entirely rewritten in accordance with the latest British
Pharmacopœia, fcap. 8vo, 7s. 6d.

III.

**NOTES ON THE ADDITIONS MADE TO THE BRITISH
PHARMACOPŒIA, 1890.** Fcap. 8vo, 1s. [Now ready.]

R. LAWTON ROBERTS, M.D. LOND., D.P.H. CAMB., M.R.C.S. ENG.
*Honorary Life Member of, and Lecturer and Examiner to the St. John Ambulance
Association.*

I.

ILLUSTRATED LECTURES ON AMBULANCE WORK.
Fourth edition, copiously Illustrated, crown 8vo, 2s. 6d. [Now ready.]

II.

**ILLUSTRATED LECTURES ON NURSING AND HY-
GIENE.** With Illustrations, crown 8vo, 2s. 6d.

ROBSON ROOSE, M.D., LL.D., F.C.S.
Fellow of the Royal College of Physicians in Edinburgh.

I.

GOUT, AND ITS RELATIONS TO DISEASES OF
 THE LIVER AND KIDNEYS. Sixth Edition, crown 8vo, 3s. 6d.

II.

NERVE PROSTRATION AND OTHER FUNCTIONAL
 DISORDERS OF DAILY LIFE. Second edition, demy 8vo, 18s.
[Just ready.]

III.

LEPROSY AND ITS TREATMENT: as Illustrated by
 Norwegian Experience. Crown 8vo, 3s. 6d.

BERNARD ROTH, F.R.C.S.

*Fellow of the Medical Society of London; Member of the Clinical and Pathological Societies
 and of the Medical Officers of Schools' Association.*

THE TREATMENT OF LATERAL CURVATURE OF
 THE SPINE. With Photographic and other Illustrations, demy 8vo,
 5s.

J. BURDON SANDERSON, M.D., LL.D., F.R.S.
Jodrell Professor of Physiology in University College, London.

UNIVERSITY COLLEGE COURSE OF PRACTICAL
 EXERCISES IN PHYSIOLOGY. With the co-operation of F. J. M.
 PAGE, B.Sc., F.C.S.; W. NORTH, B.A., F.C.S., and AUG. WALLER, M.D.
 Demy 8vo, 3s. 6d.

W. H. O. SANKEY, M.D. LOND., F.R.C.P.
Late Lecturer on Mental Diseases, University College, London, etc.

LECTURES ON MENTAL DISEASE. Second Edition, with
 coloured plates, 8vo, 12s. 6d.

JOHN SAVORY.

Member of the Society of Apothecaries, London.

A COMPENDIUM OF DOMESTIC MEDICINE AND
 COMPANION TO THE MEDICINE CHEST: Intended as a
 source of easy reference for Clergymen, Master Mariners, and Tra-
 vellers; and for Families resident at a distance from professional assist-
 ance. Tenth Edition, sm. 8vo, 5s.

E. SCHMIEGELOW, M.D.

*Consulting Physician in Laryngology to the Municipal Hospital and Director of the Oto-
 Laryngological Department in the Polyclinic at Copenhagen.*

ASTHMA: Especially in its Relation to Nasal Disease.
 Demy 8vo, 4s. 6d.

DR. B. S. SCHULTZE.

Professor of Gynecology ; Director of the Lying-in Hospital, and of the Gynecological Clinic at Jena.

THE PATHOLOGY AND TREATMENT OF DISPLACEMENTS OF THE UTERUS. Translated by J. J. MACAN, M.A., M.R.C.S. and edited by A. V. MACAN, M.B., M.Ch., Master of the Rotunda Lying-in Hospital, Dublin. With 120 Illustrations, medium 8vo, 12s. 6d.

JOHN SHAW, M.D. LOND., M.R.C.P.

Obstetric Physician to the North-West London Hospital.

ANTISEPTICS IN OBSTETRIC NURSING. A Text-book for Nurses on the Application of Antiseptics to Gynæcology and Midwifery. Coloured plate and woodcuts, 8vo, 3s. 6d.

JOHN V. SHOEMAKER, A.M., M.D.

Professor of Skin Diseases in the Medico-Chirurgical College and Hospital of Philadelphia ; Physician to the Philadelphia Hospital for Diseases of the Skin.

A PRACTICAL TREATISE ON DISEASES OF THE SKIN. Coloured Plates and other Illustrations, large 8vo, 24s.

WM. JAPP SINCLAIR, M.A., M.D.

Honorary Physician to the Manchester Southern Hospital for Women and Children, and Manchester Maternity Hospital.

ON GONORRHOËAL INFECTION IN WOMEN.
Post 8vo, 4s.

A. J. C. SKENE, M.D.

Professor of Gynecology in the Long Island College Hospital, Brooklyn, New York.

TREATISE ON THE DISEASES OF WOMEN, FOR THE USE OF STUDENTS AND PRACTITIONERS. Nine coloured plates and 251 engravings, large 8vo, 28s.

ALDER SMITH, M.B. LOND., F.R.C.S.

Resident Medical Officer, Christ's Hospital, London.

RINGWORM: Its Diagnosis and Treatment.
Third Edition, enlarged, with Illustrations, fcap. 8vo, 5s. 6d.

J. LEWIS SMITH, M.D.

Physician to the New York Infants' Hospital ; Clinical Lecturer on Diseases of Children in Bellevue Hospital Medical College.

A TREATISE ON THE DISEASES OF INFANCY AND CHILDHOOD. Fifth Edition, with Illustrations, large 8vo, 21s.

FRANCIS W. SMITH, M.B., B.S.

THE SALINE WATERS OF LEAMINGTON. Second Edit.,
with Illustrations, crown 8vo, 1s. nett.

JOHN KENT SPENDER, M.D. LOND.

Physician to the Royal Mineral Water Hospital, Bath.

THE EARLY SYMPTOMS AND THE EARLY TREATMENT OF OSTEO-ARTHRITIS, commonly called Rheumatoid Arthritis, with special reference to the Bath Thermal Waters. Sm. 8vo. 2s. 6d.

LOUIS STARR, M.D.

*Clinical Professor of Diseases of Children in the Hospital of the University of Pennsylvania ;
Physician to the Children's Hospital, Philadelphia, &c.*

HYGIENE OF THE NURSERY. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life. Second edition, with Illustrations, crown 8vo, 3s. 6d. [Just published.]

JAMES STARTIN, M.B., M.R.C.S.

Surgeon and Joint Lecturer to St. John's Hospital for Diseases of the Skin.

LECTURES ON THE PARASITIC DISEASES OF THE SKIN. VEGETOID AND ANIMAL. With Illustrations, crown 8vo, 2s. 6d.

W. R. H. STEWART, F.R.C.S., L.R.C.P. EDIN.

Aural Surgeon to the Great Northern Central Hospital; Surgeon to the London Throat Hospital, &c.

EPITOME OF DISEASES AND INJURIES OF THE EAR, with a Chapter on Naso-Pharyngeal Diseases causing Deafness. Demy 32mo, 2s. 6d.

LEWIS A. STIMSON, B.A., M.D.

Surgeon to the Presbyterian and Bellevue Hospitals; Professor of Clinical Surgery in the Medical Faculty of the University of the City of New York, &c.

A MANUAL OF OPERATIVE SURGERY.

Second Edition, with three hundred and forty-two Illustrations, post 8vo, 10s. 6d.

ADOLF STRÜMPPELL.

Director of the Medical Clinic in the University of Erlangen.

- A TEXT-BOOK OF MEDICINE FOR STUDENTS AND PRACTITIONERS.** Translated from the latest German edition by Dr. H. F. VICKERY and Dr. P. C. KNAPP, with Editorial Notes by Dr. F. C. SHATTUCK, Visiting Physician to the Massachusetts General Hospital, etc. Complete in one large vol., imp. 8vo, with 111 Illustrations, 28s.
-

JUKES DE STYRAP, M.K.Q.C.P., ETC.

Physician-Extraordinary, late Physician in Ordinary, to the Salop Infirmary; Consulting Physician to the South Salop and Montgomeryshire Infirmaries, etc.

I.

- THE YOUNG PRACTITIONER: WITH PRACTICAL HINTS AND INSTRUCTIVE SUGGESTIONS, AS SUBSIDIARY AIDS, FOR HIS GUIDANCE ON ENTERING INTO PRIVATE PRACTICE.** Demy 8vo, 7s. 6d. *nett*.

II.

- A CODE OF MEDICAL ETHICS: WITH GENERAL AND SPECIAL RULES FOR THE GUIDANCE OF THE FACULTY AND THE PUBLIC IN THE COMPLEX RELATIONS OF PROFESSIONAL LIFE.** Third edition, demy 8vo, 3s. *nett*.

III.

- MEDICO-CHIRURGICAL TARIFFS.**

Fourth Edition, fcap. 4to, revised and enlarged, 2s. *nett*.

IV.

- THE YOUNG PRACTITIONER: HIS CODE AND TARIFF.** Being the above three works in one volume. Demy 8vo, 10s. 6d. *nett*.
-

C. W. SUCKLING, M.D. LOND., M.R.C.P.

Professor of Materia Medica and Therapeutics at the Queen's College, Physician to the Queen's Hospital, Birmingham, etc.

I.

- ON THE DIAGNOSIS OF DISEASES OF THE BRAIN, SPINAL CORD, AND NERVES.** With Illustrations, crown 8vo, 8s. 6d.

II.

- ON THE TREATMENT OF DISEASES OF THE NERVOUS SYSTEM.** Crown 8vo, 7s. 6d.
-

JOHN BLAND SUTTON, F.R.C.S.

Lecturer on Comparative Anatomy, Senior Demonstrator of Anatomy, and Assistant Surgeon to the Middlesex Hospital; Erasmus Wilson Lecturer, Royal College of Surgeons, England.

- LIGAMENTS: THEIR NATURE AND MORPHOLOGY.** With numerous Illustrations, post 8vo, 4s. 6d.

HENRY R. SWANZY, A.M., M.B., F.R.C.S.I.

Examiner in Ophthalmic Surgery in the Royal University of Ireland; Surgeon to the National Eye and Ear Infirmary, Dublin; Ophthalmic Surgeon to the Adelaide Hospital, Dublin, etc.

A HANDBOOK OF THE DISEASES OF THE EYE AND THEIR TREATMENT. Third Edition, Illustrated with wood-engravings, colour tests, etc., small 8vo, 10s. 6d. [Just Published.]

EUGENE S. TALBOT, M.D., D.D.S.

Professor of Dental Surgery in the Woman's Medical College; Lecturer on Dental Pathology and Surgery in Rush Medical College, Chicago.

IRREGULARITIES OF THE TEETH AND THEIR TREATMENT. With 152 Illustrations, royal 8vo, 10s. 6d.

H. COUPLAND TAYLOR, M.D.

Fellow of the Royal Meteorological Society.

WANDERINGS IN SEARCH OF HEALTH, OR MEDICAL AND METEOROLOGICAL NOTES ON VARIOUS FOREIGN HEALTH RESORTS. Crown 8vo, with Illustrations, 6s. [Now ready.]

JOHN DAVIES THOMAS, M.D. LOND., F.R.C.S. ENG.

Physician to the Adelaide Hospital, S. Australia.

I.

HYDATID DISEASE, WITH SPECIAL REFERENCE TO ITS PREVALENCE IN AUSTRALIA. Demy 8vo, 10s. 6d.

II.

HYDATID DISEASE OF THE LUNGS. Demy 8vo, 2s.

HUGH OWEN THOMAS, M.R.C.S.

CONTRIBUTIONS TO SURGERY AND MEDICINE:—

PART I.—Intestinal Obstruction; with an Appendix on the Action of Remedies. 10s.

“ 2.—The Principles of the Treatment of Joint Disease, Inflammation, Ankylosis, Reduction of Joint Deformity, Bone Setting. 5s.

“ 3.—Fractures, Dislocations, Diseases and Deformities of the Bones of the Trunk and Upper Extremities. 10s.

“ 4.—The Collegian of 1666 and the Collegians of 1885; or what is recognised treatment? Second Edition, 1s.

“ 5.—On Fractures of the Lower Jaw. 1s.

“ 6.—The Principles of the Treatment of Fractures and Dislocations. 10s.

“ 7.—Fractures, Dislocations, Deformities, and Diseases of the Lower Extremities, 10s.

“ 8.—The Inhibition of Nerves by Drugs. Proof that Inhibitory Nerve-Fibres do not exist. 1s.

J. ASHBURTON THOMPSON, M.R.C.S.

Late Surgeon at King's Cross to the Great Northern Railway Company.

FREE PHOSPHORUS IN MEDICINE WITH SPECIAL REFERENCE TO ITS USE IN NEURALGIA. A contribution to Materia Medica and Therapeutics. An account of the History, Pharmaceutical Preparations, Dose, Internal Administration, and Therapeutic uses of Phosphorus; with a Complete Bibliography of this subject, referring to nearly 200 works upon it. Demy 8vo, 7s. 6d.

BERTRAM THORNTON, M.R.C.S., L.R.C.P. LOND.

Surgeon to the Royal Sea Bathing Infirmary for Scrofula, Margate.

THE COMPARATIVE CLIMATOLOGY OF LONDON AND THE CHIEF ENGLISH HEALTH RESORTS. With Map, crown 8vo, 1s. 6d.

J. C. THOROWGOOD, M.D.

Assistant Physician to the City of London Hospital for Diseases of the Chest.

THE CLIMATIC TREATMENT OF CONSUMPTION AND CHRONIC LUNG DISEASES. Third Edition, post 8vo, 3s. 6d.

D. HACK TUKE, M.D., LL.D.

Fellow of the Royal College of Physicians, London.

THE INSANE IN THE UNITED STATES AND CANADA. Demy 8vo, 7s. 6d.

DR. R. ULTZMANN.

ON STERILITY AND IMPOTENCE IN MAN. Translated from the German with notes and additions by ARTHUR COOPER, L.R.C.P., M.R.C.S., Surgeon to the Westminster General Dispensary. With Illustrations, fcap. 8vo, 2s. 6d.

W. H. VAN BUREN, M.D., LL.D.

Professor of Surgery in the Bellevue Hospital Medical College.

DISEASES OF THE RECTUM: And the Surgery of the Lower Bowel. Second Edition, with Illustrations, 8vo, 14s.

RUDOLPH VIRCHOW, M.D.

Professor in the University, and Member of the Academy of Sciences of Berlin, &c., &c.

INFECTION-DISEASES IN THE ARMY. Chiefly Wound Fever, Typhoid, Dysentery, and Diphtheria. Translated from the German by JOHN JAMES, M.B., F.R.C.S. Fcap. 8vo, 1s. 6d.

ALFRED VOGEL, M.D.

Professor of Clinical Medicine in the University of Dorpat, Russia.

- A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN.** Third Edition, translated and edited by H. RAPHAEL, M.D., from the Eighth German Edition, illustrated by six lithographic plates, part coloured, royal 8vo, 18s.

A. DUNBAR WALKER, M.D., C.M.

- THE PARENT'S MEDICAL NOTE BOOK.**
Oblong post 8vo, cloth, 1s. 6d.

JOHN RICHARD WARDELL, M.D. EDIN., F.R.C.P. LOND.
Late Consulting Physician to the General Hospital Tunbridge Wells.

- CONTRIBUTIONS TO PATHOLOGY AND THE PRACTICE OF MEDICINE.** Medium 8vo, 21s.

W. SPENCER WATSON, F.R.C.S. ENG., B.M. LOND.
Surgeon to the Throat Department of the Great Northern Hospital; Senior Surgeon to the Royal South London Ophthalmic Hospital.

I.

- DISEASES OF THE NOSE AND ITS ACCESSORY CAVITIES.** Second edition, with Illustrations, demy 8vo, 12s. 6d.
[Now ready.]

II.

- EYEBALL-TENSION:** Its Effects on the Sight and its Treatment. With woodcuts, p. 8vo, 2s. 6d.

III.

- ON ABSCESS AND TUMOURS OF THE ORBIT.**
Post 8vo, 2s. 6d.

FRANCIS H. WELCH, F.R.C.S.
Surgeon Major, A.M.D.

- ENTERIC FEVER:** as Illustrated by Army Data at Home and Abroad, its Prevalence and Modifications, Ætiology, Pathology and Treatment. 8vo, 5s. 6d.

W. WYNN WESTCOTT, M.B.

Deputy Coroner for Central Middlesex.

SUICIDE; its History, Literature, Jurisprudence, and Prevention. Crown 8vo, 6s.

E. G. WHITTLE, M.D. LOND., F.R.C.S. ENG.

Senior Surgeon to the Royal Alexandra Hospital for Sick Children, Brighton.

CONGESTIVE NEURASTHENIA, OR INSOMNIA AND NERVE DEPRESSION. Crown 8vo, 3s. 6d.

JOHN WILLIAMS, M.D., F.R.C.P.

Professor of Midwifery in University College, London; Obstetric Physician to University College Hospital; Physician Accoucheur to H.R.H. Princess Beatrice, &c.

CANCER OF THE UTERUS: Being the Harveian Lectures for 1886. Illustrated with Lithographic Plates, royal 8vo, 10s. 6d.

E. F. WILLOUGHBY, M.D. LOND.

THE NATURAL HISTORY OF SPECIFIC DISEASES OR STUDIES IN ETIOLOGY, IMMUNITY, AND PROPHYLAXIS. 8vo, 2s. 6d.

E. T. WILSON, B.M. OXON., F.R.C.P. LOND.

Physician to the Cheltenham General Hospital and Dispensary.

DISINFECTANTS AND HOW TO USE THEM. In Packets of one doz. price 1s.

DR. F. WINCKEL.

Formerly Professor and Director of the Gynæcological Clinic at the University of Rostock.

THE PATHOLOGY AND TREATMENT OF CHILD-BED: A Treatise for Physicians and Students. Translated from the Second German edition, with many additional notes by the Author, by J. R. CHADWICK, M.D. 8vo, 14s.

BERTRAM C. A. WINDLE, M.A., M.D. DUBL.

Professor of Anatomy in the Queen's College, Birmingham; Examiner in Anatomy in the Universities of Cambridge and Durham.

A HANDBOOK OF SURFACE ANATOMY AND LAND-MARKS. Illustrated, post 8vo, 3s. 6d.

EDWARD WOAKES, M.D. LOND.

Senior Aural Surgeon and Lecturer on Aural Surgery at the London Hospital; Surgeon to the London Throat Hospital.

I.

ON DEAFNESS, GIDDINESS AND NOISES IN THE HEAD.

VOL. I.—POST-NASAL CATARRH, AND DISEASES OF THE NOSE CAUSING DEAFNESS. With Illustrations, cr. 8vo, 6s. 6d.

VOL. II.—ON DEAFNESS, GIDDINESS AND NOISES IN THE HEAD. Third Edition, with Illustrations, cr. 8vo. [*In preparation.*]

II.

NASAL POLYPUS: WITH NEURALGIA, HAY-FEVER, AND ASTHMA, IN RELATION TO ETHMOIDITIS. With Illustrations, cr. 8vo, 4s. 6d.**DAVID YOUNG, M.C., M.B., M.D.**

Licentiate of the Royal College of Physicians, Edinburgh; Licentiate of the Royal College of Surgeons, Edinburgh, etc.

ROME IN WINTER AND THE TUSCAN HILLS IN SUMMER. A CONTRIBUTION TO THE CLIMATE OF ITALY. Small 8vo, 6s.**HERMANN VON ZEISSL, M.D.**

Late Professor at the Imperial Royal University of Vienna

OUTLINES OF THE PATHOLOGY AND TREATMENT OF SYPHILIS AND ALLIED VENEREAL DISEASES. Second Edition, revised by M. von ZEISSL, M.D., Privat-Dozent for Diseases of the Skin and Syphilis at the Imperial Royal University of Vienna. Translated, with Notes, by H. RAPHAEL, M.D., Attending Physician for Diseases of Genito-Urinary Organs and Syphilis, Bellevue Hospital, Out-Patient Department. Large 8vo, 18s.**Clinical Charts for Temperature Observations, etc.**

Arranged by W. RIGDEN, M.R.C.S. 50s. per 1000, 28s. per 500, 15s. per 250, 7s. per 100, or 1s. per dozen.

Each Chart is arranged for four weeks, and is ruled at the back for making notes of Cases; they are convenient in size, and are suitable both for hospital and private practice.

Lewis's Clinical Chart, specially designed for use with the Visiting List. This Temperature Chart is arranged for four weeks and measures 6 X 3 inches. 30s. per 1000, 16s. 6d. per 500, 3s. 6d. per 100, 1s. per 25, 6d. per 12.

Lewis's Nursing Chart.

25s. per 1000, 14s. per 500, 3s. 6d. per 100, 2s. per 50, or 1s. per 20.

These Charts afford a ready method of recording the progress of the case from day to day.

Boards to hold the Charts, price 1s.

LEWIS'S PRACTICAL SERIES.

Under this title Mr. LEWIS is publishing a Series of Monographs, embracing the various branches of Medicine and Surgery.

The volumes are written by well-known Hospital Physicians and Surgeons, recognized as authorities in the subjects of which they treat. The works are intended to be of a THOROUGHLY PRACTICAL nature, calculated to meet the requirements of the practitioner and student, and to present the most recent information in a compact and readable form.

HYGIENE AND PUBLIC HEALTH.

By LOUIS C. PARKES, M.D., D.P.H. LOND. UNIV., Fellow of the Sanitary Institute, and Member of the Board of Examiners; Assistant Professor of Hygiene and Public Health at University College, etc. Second edition, with numerous Illustrations, cr. 8vo, 9s. *[Just published.]*

MANUAL OF OPHTHALMIC PRACTICE.

By C. HIGGENS, F.R.C.S., Ophthalmic Surgeon to Guy's Hospital; Lecturer on Ophthalmology at Guy's Hospital Medical School. With Illustrations, crown 8vo, 6s.

A PRACTICAL TEXTBOOK OF THE DISEASES OF WOMEN.

By ARTHUR H. N. LEWERS, M.D. Lond., M.R.C.P. Lond., Assistant Obstetric Physician to the London Hospital; Examiner in Midwifery and Diseases of Women to the Society of Apothecaries of London, etc. Third Edition, with Illustrations, crown 8vo, 10s. 6d. *[Now ready.]*

ANÆSTHETICS THEIR USES AND ADMINISTRATION.

By DUDLEY W. BUNTON, M.D., B.S., M.R.C.P., Administrator of Anæsthetics in University College Hospital and the Hospital for Women, Soho Square. Second Edition, crown 8vo. *[In the press.]*

TREATMENT OF DISEASE IN CHILDREN: EMBODYING THE OUTLINES OF DIAGNOSIS AND THE CHIEF PATHOLOGICAL DIFFERENCES BETWEEN CHILDREN AND ADULTS.

By ANGEL MONEY, M.D., F.R.C.P., Assistant Physician to the Hospital for Sick Children, Great Ormond Street, and to University College Hospital. Second edition, cr. 8vo, 10s. 6d.

ON FEVERS: THEIR HISTORY, ETIOLOGY, DIAGNOSIS, PROGNOSIS, AND TREATMENT.

By ALEXANDER COLLIE, M.D. Aberd., Member of the Royal College of Physicians of London; Medical Superintendent of the Eastern Hospitals; Secretary of the Epidemiological Society for Germany and Russia. Illustrated with Coloured Plates, crown 8vo, 8s. 6d.

HANDBOOK OF DISEASES OF THE EAR FOR THE USE OF STUDENTS AND PRACTITIONERS.

By URBAN PRITCHARD, M.D. Edin., F.R.C.S. Eng., Professor of Aural Surgery at King's College, London; Aural Surgeon to King's College Hospital; Senior Surgeon to the Royal Ear Hospital. Second Edition, with Illustrations, crown 8vo, 5s. *[Now ready.]*

A PRACTICAL TREATISE ON DISEASES OF THE KIDNEYS AND URINARY DERANGEMENTS.

By CHARLES HENRY RALFE, M.A., M.D. Cantab., Fellow of the Royal College of Physicians, London; Assistant Physician to the London Hospital; Examiner in Medicine to the University of Durham, etc., etc. With Illustrations, crown 8vo, 10s. 6d.

DENTAL SURGERY FOR MEDICAL PRACTITIONERS AND STUDENTS OF MEDICINE.

By ASHLEY W. BARRETT, M.B. Lond., M.R.C.S., L.S.D., Dental Surgeon to, and Lecturer on Dental Surgery in the Medical School of, the London Hospital. Second edition, with Illustrations, cr. 8vo, 3s. 6d.

BODILY DEFORMITIES AND THEIR TREATMENT: A HANDBOOK OF PRACTICAL ORTHOPÆDICS.

By H. A. REEVES, F.R.C.S. Edin., Senior Assistant Surgeon and Teacher of Practical Surgery at the London Hospital; Surgeon to the Royal Orthopædic Hospital, &c. With numerous Illustrations, cr. 8vo, 8s. 6d.

Further volumes will be announced in due course.

THE NEW SYDENHAM SOCIETY'S PUBLICATIONS.

President :—SIR JAMES PAGET, BART., F.R.S.*Honorary Secretary* :—JONATHAN HUTCHINSON, ESQ., F.R.S.*Treasurer* :—W. SEDGWICK SAUNDERS, M.D., F.S.A.

Annual Subscription, One Guinea.

The Society issues translations of recent standard works by continental authors on subjects of general interest to the profession.

Amongst works recently issued are "Flügge's Micro-Organisms," "Cohnheim's Pathology," "Henoch's Children," "Spiegelberg's Midwifery," "Hirsch's Historical and Geographical Pathology," "Essays on Micro-Parasites," works by Charcot, Duchenne, Begbie, Billroth, Graves, Koch, Hebra, Guttman, etc.

The Society also has in hand an Atlas of Pathology with Coloured Plates, and a valuable and exhaustive "Lexicon of Medicine and the Allied Sciences."

The Annual Report, with full list of works published, and all further information will be sent on application.

PERIODICAL WORKS PUBLISHED BY H. K. LEWIS.

THE BRITISH JOURNAL OF DERMATOLOGY. Edited by H. G. Brooke, H. Radcliffe Crocker, T. Colcott Fox, Malcolm Morris, J. F. Payne and J. J. Pringle. Published monthly, 1s. Annual Subscription 12s. post free.

THE NEW YORK MEDICAL JOURNAL. A Weekly Review of Medicine. Annual Subscription, Thirty Shillings, post free.

THE THERAPEUTIC GAZETTE. A Monthly Journal, devoted to the Science of Pharmacology, and to the introduction of New Therapeutic Agents. Edited by Dr. R. M. Smith. Annual Subscription, 10s., post free.

THE GLASGOW MEDICAL JOURNAL. Published Monthly. Annual Subscription 20s., post free. Single numbers, 2s. each.

LIVERPOOL MEDICO-CHIRURGICAL JOURNAL, including the Proceedings of the Liverpool Medical Institution. Published twice yearly, 3s. 6d. each number.

TRANSACTIONS OF THE COLLEGE OF PHYSICIANS OF PHILADELPHIA. Volumes I. to VI., 8vo, 10s. 6d. each.

MIDDLESEX HOSPITAL, REPORTS OF THE MEDICAL, SURGICAL, AND Pathological Registrars for 1883 to 1888. Demy 8vo, 2s. 6d. *net* each volume.

* * MR. LEWIS is in constant communication with the leading publishing firms in America, and has transactions with them for the sale of his publications in that country. Advantageous arrangements are made in the interests of Authors for the publishing of their works in the United States.

Mr. Lewis's publications can be procured of all Booksellers in any part of the world.

24

